

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 November 2000 (30.11.2000)

PCT

(10) International Publication Number
WO 00/71535 A1


- (51) International Patent Classification⁷: C07D 401/06, A61K 31/404, A61P 43/00, C07D 209/24, 401/14
- (21) International Application Number: PCT/US00/14003
- (22) International Filing Date: 19 May 2000 (19.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/316,761 21 May 1999 (21.05.1999) US
60/154,594 17 September 1999 (17.09.1999) US
60/202,608 9 May 2000 (09.05.2000) US
- (71) Applicant: SCIOS INC. [US/US]; 820 West Maude Avenue, Sunnyvale, CA 94087 (US).
- (72) Inventors: MAVUNKEL, Babu, J.; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). CHAKRAVARTY, Sarvajit; 976-2 Alpine Terrace, Sunnyvale, CA 94086 (US). PERUMATTAM, John, J.; 30 Chester Circle, Los Altos, CA 94022 (US). DUGAR, Sundeep; 5943 Sterling Oaks Drive, San Jose, CA 95120 (US). LU, Qing; 350 Foresail Court, Foster City, CA 94404 (US). LIANG, Xi; 1011 Rosa Avenue, Sunnyvale, CA 94086 (US).
- (74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Forster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).
- (81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

[Continued on next page]

(54) Title: INDOLE-TYPE DERIVATIVES AS INHIBITORS OF p38 KINASE



(57) Abstract: The invention is directed to methods to inhibit p38- α kinase using compounds of formula (1), and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein (a) represents a single or double bond; one Z² is CA or CR⁸A and the other is CR¹, CR¹₂, NR⁶ or N wherein each R¹, R⁶ and R⁸ is independently hydrogen or noninterfering substituent; A is -W-COX_iY wherein Y is COR² or an isostere thereof and R² is hydrogen or a noninterfering substituent, each of W and X is a spacer of 2-6Å, and each of i and j is independently 0 or 1; Z³ is NR⁷ or O; each R³ is independently a noninterfering substituent; n is 0-3; each of L¹ and L² is a linker; each R⁴ is independently a noninterfering substituent; m is 0-4; Z¹ is CR⁵ or N wherein R⁵ is hydrogen or a noninterfering substituent; each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3; Ar is an aryl group substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; and the distance between the atom of Ar linked to L² and the center of the α ring is 4.5-24Å.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INDOLE-TYPE DERIVATIVES AS INHIBITORS OF p38 KINASE

This application claims priority under 35 U.S.C. § 119(e) to U.S. Serial No. (Attorney Docket No. 21900-30290.20) filed 9 May 2000 and to U.S. Serial No. 60/154,594 filed 17 September 1999. Priority is claimed under 35 U.S.C. § 120 with respect to U.S. Serial No. 09/316,761 filed 21 May 1999. The contents of these applications are incorporated herein by reference in their entirety.

Field of the Invention

The invention relates to treating various disorders associated with enhanced activity of kinase p38- α . More specifically, it concerns compounds that are related to indole-type derivatives coupled to piperazine- or piperidine-type moieties as useful in these methods.

Background Art

A large number of chronic and acute conditions have been recognized to be associated with perturbation of the inflammatory response. A large number of cytokines participate in this response, including IL-1, IL-6, IL-8 and TNF. It appears that the activity of these cytokines in the regulation of inflammation rely at least in part on the activation of an enzyme on the cell signaling pathway, a member of the MAP kinase family generally known as p38 and alternatively known as CSBP and RK. This kinase is activated by dual phosphorylation after stimulation by physiochemical stress, treatment with lipopolysaccharides or with proinflammatory cytokines such as IL-1 and TNF. Therefore, inhibitors of the kinase activity of p38 are useful anti-inflammatory agents.

Eye diseases associated with a fibroproliferative condition include retinal reattachment surgery accompanying proliferative vitreoretinopathy, cataract extraction with intraocular lens implantation, and post glaucoma drainage surgery.

PCT applications WO98/06715, WO98/07425, and WO 96/40143, all of which are incorporated herein by reference, describe the relationship of p38 kinase inhibitors with various disease states. As mentioned in these applications, inhibitors of p38 kinase are useful in treating a variety of diseases associated with chronic inflammation. These applications list rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, Gram-negative sepsis,

toxic shock syndrome, asthma, adult respiratory distress syndrome, stroke, reperfusion injury, CNS injuries such as neural trauma and ischemia, psoriasis, restenosis, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases such as osteoporosis, graft-versus-host reaction, Crohn's Disease, ulcerative colitis including inflammatory bowel disease (IBD) and pyresis.

The above-referenced PCT applications disclose compounds which are p38 kinase inhibitors said to be useful in treating these disease states. These compounds are either imidazoles or are indoles substituted at the 3- or 4-position with a piperazine ring linked through a carboxamide linkage. Additional compounds which are conjugates of piperazines with indoles are described as insecticides in WO97/26252, also incorporated herein by reference.


Certain aroyl/phenyl-substituted piperazines and piperidines which inhibit p38- α kinase are described in PCT publication WO00/12074 published 9 March 2000. In addition, indolyl substituted piperidines and piperazines which inhibit this enzyme are described in PCT publication No. WO99/61426 published 2 December 1999. Carbolene derivatives of piperidine and piperazine as p38- α inhibitors are described in PCT/US00/07934 filed 24 March 2000.

None of the foregoing patents describes the indole derivatives described herein which specifically inhibit p38- α .


Disclosure of the Invention

The invention is directed to methods and compounds useful in treating conditions that are characterized by enhanced p38- α activity. These conditions include inflammation, proliferative diseases, and certain cardiovascular disorders as well as Alzheimer's disease as further described below.

Compounds of the invention have been found to inhibit p38 kinase, the α -isoform in particular, and are thus useful in treating diseases mediated by these activities. The compounds of the invention are of the formula



and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein

 represents a single or double bond;

one Z² is CA or CR⁸A and the other is CR¹, CR¹₂, NR⁶ or N wherein each R¹, R⁶
5 and R⁸ is independently hydrogen or noninterfering substituent;

A is -W_i-COX_jY wherein Y is COR² or an isostere thereof and R² is hydrogen or a noninterfering substituent, each of W and X is a spacer of 2-6Å, and each of i and j is independently 0 or 1;

Z³ is NR⁷ or O;

10 each R³ is independently a noninterfering substituent;

n is 0-3;

each of L¹ and L² is a linker;

each R⁴ is independently a noninterfering substituent;

m is 0-4;

15 Z¹ is CR⁵ or N wherein R⁵ is hydrogen or a noninterfering substituent;

each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3;

Ar is an aryl group substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; and

the distance between the atom of Ar linked to L² and the center of the α ring is 4.5-24Å.

20 The invention is directed to methods of treating inflammation or proliferative conditions using these compounds. The invention is also directed to treating conditions associated with cardiac failure and Alzheimer's disease using the invention compounds.

Modes of Carrying Out the Invention

The compounds of formula (1) are useful in treating conditions which are characterized by
25 overactivity of p38 kinase, in particular the α-isoform. Conditions "characterized by enhanced p38-α activity" include those where this enzyme is present in increased amount or wherein the enzyme has been modified to increase its inherent activity, or both. Thus, "enhanced activity" refers to any condition wherein the effectiveness of these proteins is undesirably high, regardless of the cause.

30 The compounds of the invention are useful in conditions where p38-α kinase shows enhanced activity. These conditions are those in which fibrosis and organ sclerosis are

caused by, or accompanied by, inflammation, oxidation injury, hypoxia, altered temperature or extracellular osmolarity, conditions causing cellular stress, apoptosis or necrosis. These conditions include ischemia-reperfusion injury, congestive heart failure, progressive pulmonary and bronchial fibrosis, hepatitis, arthritis, inflammatory bowel disease, glomerular sclerosis, interstitial renal fibrosis, chronic scarring diseases of the eyes, bladder and reproductive tract, bone marrow dysplasia, chronic infectious or autoimmune states and traumatic or surgical wounds. These conditions, of course, would be benefited by compounds which inhibit p38- α . Methods of treatment with the compounds of the invention are further discussed below.

The Invention Compounds

The compounds useful in the invention are derivatives of indole-type compounds containing a mandatory substituent, A, at a position corresponding to the 2- or 3- position of indole. In general, an indole-type nucleus is preferred, although alternatives within the scope of the invention are also illustrated below.

In the description above, certain positions of the molecule are described as permitting "noninterfering substituents." This terminology is used because the substituents in these positions generally speaking are not relevant to the essential activity of the molecule taken as a whole. A wide variety of substituents can be employed in these positions, and it is well within ordinary skill to determine whether any particular arbitrary substituent is or is not "noninterfering."

As used herein, a "noninterfering substituent" is a substituent which leaves the ability of the compound of formula (1) to inhibit p38- α activity qualitatively intact. Thus, the substituent may alter the degree of inhibition of p38- α . However, as long as the compound of formula (1) retains the ability to inhibit p38- α activity, the substituent will be classified as "noninterfering." A number of assays for determining the ability of any compound to inhibit p38- α activity are available in the art. A whole blood assay for this evaluation is illustrated below: the gene for p38- α has been cloned and the protein can be prepared recombinantly and its activity assessed, including an assessment of the ability of an arbitrarily chosen compound to interfere with this activity. The essential features of the molecule are tightly defined. The positions which are occupied by "noninterfering substituents" can be substituted by conventional organic moieties as is understood in the

art. It is irrelevant to the present invention to test the outer limits of such substitutions.

The essential features of the compounds are those set forth with particularity herein.

In addition, L¹ and L² are described herein as linkers. The nature of such linkers is less important that the distance they impart between the portions of the molecule. Typical

5 linkers include alkylene, *i.e.* (CH₂)_n-R; alkenylene - *i.e.*, an alkylene moiety which contains a double bond, including a double bond at one terminus. Other suitable linkers include, for example, substituted alkylenes or alkenylenes, carbonyl moieties, and the like. As used herein, "hydrocarbyl residue" refers to a residue which contains only carbon and hydrogen. The residue may be aliphatic or aromatic, straight-chain, cyclic, branched,
10 saturated or unsaturated. The hydrocarbyl residue, when so stated however, may contain heteroatoms over and above the carbon and hydrogen members of the substituent residue. Thus, when specifically noted as containing such heteroatoms, the hydrocarbyl residue may also contain carbonyl groups, amino groups, hydroxyl groups and the like, or contain heteroatoms within the "backbone" of the hydrocarbyl residue.

15 As used herein, "inorganic residue" refers to a residue that does not contain carbon.

Examples include, but are not limited to, halo, hydroxy, NO₂ or NH₂.

As used herein, the term "alkyl," "alkenyl" and "alkynyl" include straight- and branched-chain and cyclic monovalent substituents. Examples include methyl, ethyl, isobutyl, cyclohexyl, cyclopentylethyl, 2-propenyl, 3-butenyl, and the like. Typically, the alkyl,
20 alkenyl and alkynyl substituents contain 1-10C (alkyl) or 2-10C (alkenyl or alkynyl). Preferably they contain 1-6C (alkyl) or 2-6C (alkenyl or alkynyl). Heteroalkyl, heteroalkenyl and heteroalkynyl are similarly defined but may contain 1-2 O, S or N heteroatoms or combinations thereof within the backbone residue.

As used herein, "acyl" encompasses the definitions of alkyl, alkenyl, alkynyl and the
25 related hetero-forms which are coupled to an additional residue through a carbonyl group.

"Aromatic" moiety refers to a monocyclic or fused bicyclic moiety such as phenyl or naphthyl; "heteroaromatic" also refers to monocyclic or fused bicyclic ring systems containing one or more heteroatoms selected from O, S and N. The inclusion of a heteroatom permits inclusion of 5-membered rings as well as 6-membered rings. Thus,
30 typical aromatic systems include pyridyl, pyrimidyl, indolyl, benzimidazolyl, benzotriazolyl, isoquinolyl, quinolyl, benzothiazolyl, benzofuranyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl and the like. Any monocyclic or fused ring bicyclic system

which has the characteristics of aromaticity in terms of electron distribution throughout the ring system is included in this definition. Typically, the ring systems contain 5-12 ring member atoms.

Similarly, "arylalkyl" and "heteroalkyl" refer to aromatic and heteroaromatic systems which are coupled to another residue through a carbon chain, including substituted or unsubstituted, saturated or unsaturated, carbon chains, typically of 1-6C. These carbon chains may also include a carbonyl group, thus making them able to provide substituents as an acyl moiety.

When the compounds of Formula 1 contain one or more chiral centers, the invention

includes optically pure forms as well as mixtures of stereoisomers or enantiomers

With respect to the portion of the compound between the atom of Ar bound to L² and ring α , L¹ and L² are linkers which space the substituent Ar from ring α at a distance of 4.5-24Å, preferably 6-20Å, more preferably 7.5-10Å. The distance is measured from the

center of the α ring to the atom of Ar to which the linker L² is attached. Typical, but

nonlimiting, embodiments of L¹ and L² are CO and isosteres thereof, or optionally

substituted isosteres, or longer chain forms. L², in particular, may be alkylene or

alkenylene optionally substituted with noninterfering substituents or L¹ or L² may be or

may include a heteroatom such as N, S or O. Such substituents include, but are limited to, a moiety selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl,

aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl,

halo, OR, NR₂, SR, SOR, SO₂R, OCOR, NRCOR, NRCONR₂, NRCOOR, OCONR₂, RCO, COOR, alkyl-OOR, SO₃R, CONR₂, SO₂NR₂, NRSO₂NR₂, CN, CF₃, R₃Si, and NO₂,

wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof, and

wherein two substituents on L² can be joined to form a non-aromatic saturated or

unsaturated ring that includes 0-3 heteroatoms which are O, S and/or N and which contains 3 to 8 members or said two substituents can be joined to form a carbonyl moiety or an oxime, oximeether, oximeester or ketal of said carbonyl moiety.

Isosteres of CO and CH₂, include SO, SO₂, or CHOH. CO and CH₂ are preferred.


Thus, L² is substituted with 0-2 substituents. Where appropriate, two optional substituents

on L² can be joined to form a non-aromatic saturated or unsaturated hydrocarbonyl ring that includes 0-3 heteroatoms such as O, S and/or N and which contains 3 to 8 members. Two

optional substituents on L² can be joined to form a carbonyl moiety which can be subsequently converted to an oxime, an oximeether, an oximeester, or a ketal.

Ar is aryl, heteroaryl, including 6-5 fused heteroaryl, cycloaliphatic or cycloheteroaliphatic that can be optionally substituted. Ar is preferably optionally substituted phenyl.

- 5 Each substituent on Ar is independently a hydrocarbyl residue (1-20C) containing 0-5 heteroatoms selected from O, S and N, or is an inorganic residue. Preferred substituents include those selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR₂, SR, SOR, SO₂R, OCOR, NRCOR, NRCONR₂, NR₂COOR, 10 OCONR₂, RCO, COOR, alkyl-OOR, SO₃R, CONR₂, SO₂NR₂, NRSO₂NR₂, CN, CF₃, R₃Si, and NO₂, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof, and wherein two of said optional substituents on adjacent positions can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members. More preferred substituents include halo, alkyl (1-4C) 15 and more preferably, fluoro, chloro and methyl. These substituents may occupy all available positions of the aryl ring of Ar, preferably 1-2 positions, most preferably one position. These substituents may be optionally substituted with substituents similar to those listed. Of course some substituents, such as halo, are not further substituted, as known to one skilled in the art.
- 20 Two substituents on Ar can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members.
- Between L¹ and L² is a piperidine-type moiety of the following formula:



- 25 Z¹ is CR⁵ or N wherein R⁵ is H or a noninterfering substituent. Each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3. The noninterfering substituents R⁵ include, without limitation, halo, alkyl, alkoxy, aryl, arylalkyl, aryloxy, heteroaryl, acyl, carboxy, or hydroxy. Preferably, R⁵ is H, alkyl, OR, NR₂, SR or halo, where R is H or alkyl. Additionally, R⁵ can be joined with an R⁴ substituent to form an optionally substituted non-aromatic saturated or unsaturated hydrocarbyl ring which contains 3-8

members and 0-3 heteroatoms such as O, N and/or S. Preferred embodiments include compounds wherein Z^1 is CH or N, and those wherein both l and k are 1.

R^4 represents a noninterfering substituent such as a hydrocarbyl residue (1-20C)

containing 0-5 heteroatoms selected from O, S and N. Preferably R^4 is alkyl, alkoxy, aryl,

5 arylalkyl, aryloxy, heteroalkyl, heteroaryl, heteroarylalkyl, RCO, =O, acyl, halo, CN, OR, NRCOR, NR, wherein R is H, alkyl (preferably 1-4C), aryl, or hetero forms thereof. Each

appropriate substituent is itself unsubstituted or substituted with 1-3 substituents. The

substituents are preferably independently selected from a group that includes alkyl,

alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl,

10 heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR_2 , SR, SOR, SO_2R , OCOR,

NRCOR, $NRCONR_2$, $NRCOOR$, $CONR_2$, RCO, COOR, alkyl-OOR, SO_3R , $CONR_2$,

SO_2NR_2 , $NRSO_2NR_2$, CN, CF_3 , R_3Si , and NO_2 , wherein each R is independently H, alkyl,

alkenyl or aryl or heteroforms thereof and two of R^4 on adjacent positions can be joined to

form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring

15 which contains 3-8 members, or R^4 is =O or an oxime, oximeether, oximeester or ketal

thereof. R^4 may occur m times on the ring; m is an integer of 0-4. Preferred embodiments

of R^4 comprise alkyl (1-4C) especially two alkyl substituents and carbonyl. Most

preferably R^4 comprises two methyl groups at positions 2 and 5 or 3 and 6 of a piperidinyl

or piperazinyl ring or =O preferably at the 5-position of the ring. The substituted forms

20 may be chiral and an isolated enantiomer may be preferred.

R^3 also represents a noninterfering substituent. Such substituents include hydrocarbyl

residues (1-6C) containing 0-2 heteroatoms selected from O, S and/or N and inorganic

residues. n is an integer of 0-3, preferably 0 or 1. Preferably, the substituents represented

by R^3 are independently halo, alkyl, heteroalkyl, OCOR, OR, NRCOR, SR, or NR_2 ,

25 wherein R is H, alkyl, aryl, or heteroforms thereof.. More preferably R^3 substituents are

selected from alkyl, alkoxy or halo, and most preferably methoxy, methyl, and chloro.

Most preferably, n is 0 and the α ring is unsubstituted, except for L^1 or n is 1 and R^3 is halo

or methoxy.

In the ring labeled β , Z^3 may be NR^7 or O - i.e., the compounds may be related to indole or

30 benzofuran. If C^3 is NR^7 , preferred embodiments of R^7 include H or optionally substituted

alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl,

heteroalkynyl, heteroalkylaryl, or is SOR, SO_2R , RCO, COOR, alkyl-COR, SO_3R ,

CONR₂, SO₂NR₂, CN, CF₃, NR₂, OR, alkyl-SR, alkyl-SOR, alkyl-SO₂R, alkyl-OCOR, alkyl-COOR, alkyl-CN, alkyl-CONR₂, or R₃Si, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof. More preferably, R⁷ is hydrogen or is alkyl (1-4C), preferably methyl or is acyl (1-4C), or is COOR wherein R is H, alkyl, alkenyl or aryl or hetero forms thereof. R⁷ is also preferably a substituted alkyl wherein the preferred substituents are form ether linkages or contain sulfinic or sulfonic acid moieties. Other preferred substituents include sulfhydryl substituted alkyl substituents. Still other preferred substituents include CONR₂ wherein R is defined as above.

It is preferred that the indicated dotted line represents a double bond; however, compounds which contain a saturated β ring are also included within the scope of the invention.

Preferably, the mandatory substituent CA or CR⁸A is in the 3- position; regardless of which position this substituent occupies, the other position is CR¹, CR¹₂, NR⁶ or N. CR¹ is preferred. Preferred embodiments of R¹ include hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl,

heteroalkylaryl, NH-aroyl, halo, OR, NR₂, SR, SOR, SO₂R, OCOR, NRCOR, NRCONR₂, NRCOOR, OCONR₂, RCO, COOR, alkyl-OOR, SO₃R, CONR₂, SO₂NR₂, NRSO₂NR₂, CN, CF₃, R₃Si, and NO₂, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof and two of R¹ can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members. Most preferably, R¹ is H, alkyl, such as methyl, most preferably, the ring labeled α contains a double bond and CR¹ is CH or C-alkyl. Other preferable forms of R¹ include H, alkyl, acyl, aryl, arylalkyl, heteroalkyl, heteroaryl, halo, OR, NR₂, SR, NRCOR, alkyl-OOR, RCO, COOR, and CN, wherein each R is independently H, alkyl, or aryl or heteroforms thereof.

While the position not occupied by CA is preferred to include CR¹, the position can also be N or NR⁶. While NR⁶ is less preferred (as in that case the ring labeled β would be saturated), if NR⁶ is present, preferred embodiments of R⁶ include H, or alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, or is SOR, SO₂R, RCO, COOR, alkyl-COR, SO₃R, CONR₂, SO₂NR₂, CN, CF₃, or R₃Si wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof.

Preferably, CR⁸A or CA occupy position 3- and preferably Z² in that position is CA. However, if the β ring is saturated and R⁸ is present, preferred embodiments for R⁸ include H, halo, alkyl, alkenyl and the like. Preferably R⁸ is a relatively small substituent corresponding, for example, to H or lower alkyl 1-4C.

- 5 A is -W_i-COX_jY wherein Y is COR² or an isostere thereof and R² is a noninterfering substituent. Each of W and X is a spacer and may be, for example, optionally substituted alkyl, alkenyl, or alkynyl, each of i and j is 0 or 1. Preferably, W and X are unsubstituted. Preferably, j is 0 so that the two carbonyl groups are adjacent to each other. Preferably, also, i is 0 so that the proximal CO is adjacent the ring. However, compounds wherein the
- 10 proximal CO is spaced from the ring can readily be prepared by selective reduction of an initially glyoxal substituted β ring. In the most preferred embodiments of the invention, the α/β ring system is an indole containing CA in position 3- and wherein A is COCR². The noninterfering substituent represented by R², when R² is other than H, is a hydrocarbyl residue (1-20C) containing 0-5 heteroatoms selected from O, S and/or N or is
- 15 an inorganic residue. Preferred are embodiments wherein R² is H, or is straight or branched chain alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroalkyl, heteroaryl, or heteroarylalkyl, each optionally substituted with halo, alkyl, heteroalkyl, SR, OR, NR₂, OCOR, NRCOR, NRCONR₂, NRSO₂R, NRSO₂NR₂, OCONR₂, CN, COOR, CONR₂, COR, or R₃Si wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-
- 20 containing forms thereof, or wherein R² is OR, NR₂, SR, NRCONR₂, OCONR₂, or NRSO₂NR₂, wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof, and wherein two R attached to the same atom may form a 3-8 member ring and wherein said ring may further be substituted by alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroalkyl, heteroaryl, heteroarylalkyl, each optionally substituted with
- 25 halo, SR, OR, NR₂, OCOR, NRCOR, NRCONR₂, NRSO₂R, NRSO₂NR₂, OCONR₂, or R₃Si wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof wherein two R attached to the same atom may form a 3-8 member ring, optionally substituted as above defined.
- Other preferred embodiments of R² are H, heteroarylalkyl, -NR₂, heteroaryl, -COOR, -NHRNR₂, heteroaryl-COOR, heteroaryloxy, -OR, heteroaryl-NR₂, -NROR and alkyl.
- 30 Most preferably R² is isopropyl piperazinyl, methyl piperazinyl, dimethylamine, piperazinyl, isobutyl carboxylate, oxycarbonylethyl, morpholinyl,

aminoethyldimethylamine, isobutyl carboxylate piperazinyl, oxypiperazinyl, ethylcarboxylate piperazinyl, methoxy, ethoxy, hydroxy, methyl, amine, aminoethyl pyrrolidinyl, aminopropanediol, piperidinyl, pyrrolidinyl-piperidinyl, or methyl piperidinyl.

- 5 Isosteres of COR² as represented by Y are defined as follows.

The isosteres have varying lipophilicity and may contribute to enhanced metabolic stability. Thus, Y, as shown, may be replaced by the isosteres in Table 1.





Table 1 - Acid Isosteres		
Names of Groups	Chemical Structures	Substitution Groups (SG)
tetrazole		n/a
1,2,3-triazole		H; SCH ₃ ; COCH ₃ ; Br; SOCH ₃ ; SO ₂ CH ₃ ; NO ₂ ; CF ₃ ; CN; COOMe
1,2,4-triazole		H; SCH ₃ ; COCH ₃ ; Br; SOCH ₃ ; SO ₂ CH ₃ ; NO ₂
imidazole		H; SCH ₃ ; COCH ₃ ; Br; SOCH ₃ ; SO ₂ CH ₃ ; NO ₂

Thus, isosteres include tetrazole, 1,2,3-triazole, 1,2,4-triazole and imidazole.

The compounds of formula (1) may be supplied in the form of their pharmaceutically acceptable acid-addition salts including salts of inorganic acids such as hydrochloric, sulfuric, hydrobromic, or phosphoric acid or salts of organic acids such as acetic, tartaric, succinic, benzoic, salicylic, and the like. If a carboxyl moiety is present on the compound of formula (1), the compound may also be supplied as a salt with a pharmaceutically acceptable cation.


Synthesis of the Invention Compounds

The following Reaction Scheme is illustrative of the conversion of a 4-benzyl piperidinyl-indole-5-carboxamide to the glyoxalic acid compounds of the invention and derivatives thereof.




$\text{X} = \text{OCH}_3, \text{Cl}, \text{CH}_3$; $\text{Y} = \text{H}, \text{Halogen etc.}$

$\text{NR}_1\text{R}_2 = \text{NH}_2, \text{NH-Alkyl}, \text{NH-Aryl}, \text{N-Dialkyl},$




Of course, the 4-benzyl piperidinyl carbonyl of the illustration at position 5 may be generalized as




and the glyoxal type substituent at position 3 can be generalized to $W_i\text{COX}_j\text{Y}$.

5 Similarly, embodiments wherein the indole-type moiety is




can be used in these schemes. Methods to synthesize the compounds of the invention are, in general, known in the art.

The following general schemes illustrate such methods.




Scheme 1

Substituted amino benzoic acid esters such as **I** can be treated with reagents such as thiomethylacetaldehyde dimethyl acetal and N-chlorosuccinamide in methylene chloride at low temperature followed by the treatment with a base such as triethylamine at reflux in methylene chloride, dichloroethane or chloroform to give indoles **II**, Scheme 1. Treatment with reagents such as Raney-Nickel in an appropriate solvent such as ethanol, methanol or isopropanol will yield the corresponding indole carboxylic acid ester which when hydrolyzed under base conditions will give the desired substituted indole carboxylic acid.




Scheme 2

Alternatively, substituted amino benzoic acid esters **I** can be converted to the ketals **IV**, Scheme 2, with an appropriate aldehyde under conditions of reductive alkylation with reagents such as sodium triacetoxyborohydride in acetic acid in the presence of sodium sulfate. The amines can then be treated with lewis acids such as aluminum chloride, titanium chloride, BF_3 -etherate in dichloromethane or dichloroethane, under reflux to give the corresponding substituted indole methyl esters, with appropriate substitutions.



Scheme 3


Another method could involve the treatment of the substituted amino benzoic acid esters **I** with iodine and sodium periodate in an appropriate solvent such as dimethylformamide, to give the corresponding iodo aniline **V**, Scheme 3. This can be coupled with an acetylene such as trimethyl silyl acetylene or ethylethynyl ether in the presence of an appropriate catalysts such as palladium and copper and a base such as triethylamine to give the silyl coupled product such as **VI**. Subsequent cyclization in a solvent such as dimethylformamide and in the presence of a catalyst such as copper iodide would give the appropriately substituted indoles **VII**.



Scheme 4


Synthesis of the required piperidines can be achieved by treating an appropriate piperidone such as **VIII**, Scheme 4, with substituted benzyl phosphonate esters in the presence of a

base such as sodium hydride to give alkenes which can be reduced to the corresponding substituted 4-benzylpiperidine such as **IX**. The hydrogenations are typically done in the presence of catalytic metals in solvents such as methanol, ethanol and ethyl acetate.




Scheme 5

An alternate method could involve isonipecotoyl chlorides such as **X** which can be used to acylate appropriately substituted benzenes (ArH) in the presence of a lewis acid such as aluminum chloride to give the ketones **XI**, Scheme 5. Further modifications of the carbonyl moiety of **XI** using methods and routes generally known can then lead to the desired compounds **XII**.




Scheme 6

Substituted piperazines can be reacted with various and appropriate ArL^2X in the presence or absence of a base or other catalytic reagent to give the substituted piperazines **XV**, Scheme 6. These can be further resolved to the chiral components with the use a chiral resolving agent such as tartaric acid to give either enantiomers of the substituted piperazines **XV**.




Scheme 7

Compounds **III** can be treated with halides, acid chlorides and other electrophiles (BX), Scheme 7, containing a variety of different substituents, in the presence of a base such as sodium hydride, in a variety of different solvents, to give compounds of type **XVI**. These can then be converted to the corresponding acids **XVII** by treatment with appropriate reagents such as an aqueous base. The acids are then coupled to substituted amines **IX**, **XII** or **XV** using a coupling agent such as EDAC.HCl in a variety of solvents including methylene chloride, dimethyl formamide, to give compounds **XVIII**.



Scheme 8

Compounds **XVIII** can be first treated with acid chlorides such as oxalyl chloride in methylene chloride under anhydrous conditions followed by treatment with a variety of nucleophiles WH to give compounds of type **XIX**, Scheme 8.



Scheme 9

Compounds of type **XXIV** can be synthesized starting with the appropriately substituted benzofurans of type **XX** and coupling them with amines **IX**, **XII** or **XV** in the presence of relevant coupling reagents to give compounds **XXI**, Scheme 9. Subsequent acylation to **XXII** can be achieved with an acylating agent such as acetic anhydride in the presence of a catalyst such as Fe(III). Oxidation of the acetyl moiety of **XXII** to the glyoxalic acid moiety of **XXIII** can be accomplished using an oxidizing agent such as selenium dioxide (Ref: F Da Settimo et al. Eur. J. Med. Chem (1996), 31, 951-956; M. C. Cook, et al. (1975) Br patent 1, 399,089.; E. Campaigne, et al. J. Med. Chem. (1965), 136-137). Finally coupling of the acid with the appropriate nucleophile in WH can be achieved using any one of the variety of coupling agents known in a variety of solvents to give compounds of type **XXIV**.

Assays for p38 α Kinase Inhibition

For each of the assay procedures described below, the TNF- α production correlates to the activity of p38- α kinase.

A. Human Whole Blood Assay for p38 Kinase Inhibition

Venous blood is collected from healthy male volunteers into a heparinized syringe and is used within 2 hours of collection. Test compounds are dissolved in 100% DMSO and 1 μ l aliquots of drug concentrations ranging from 0 to 1 mM are dispensed into quadruplicate wells of a 24-well microtiter plate (Nunclon Delta SI, Applied Scientific, So. San Francisco, CA). Whole blood is added at a volume of 1 ml/well and the mixture is incubated for 15 minutes with constant shaking (Titer Plate Shaker, Lab-Line Instruments, Inc., Melrose Park, IL) at a humidified atmosphere of 5% CO₂ at 37°C. Whole blood is cultured either undiluted or at a final dilution of 1:10 with RPMI 1640 (Gibco 31800 + NaHCO₃, Life Technologies, Rockville, MD and Scios, Inc., Sunnyvale, CA). At the end of the incubation period, 10 μ l of LPS (*E. coli* 0111:B4, Sigma Chemical Co., St. Louis, MO) is added to each well to a final concentration of 1 or 0.1 μ g/ml for undiluted or 1:10 diluted whole blood, respectively. The incubation is continued for an additional 2 hours. The reaction is stopped by placing the microtiter plates in an ice bath and plasma or cell-free supernates are collected by centrifugation at 3000 rpm for 10 minutes at 4°C. The

plasma samples are stored at -80°C until assayed for TNF- α levels by ELISA, following the directions supplied by Quantikine Human TNF- α assay kit (R&D Systems, Minneapolis, MN).

IC₅₀ values are calculated using the concentration of inhibitor that causes a 50% decrease as compared to a control.

B. Enriched Mononuclear Cell Assay for p38 Kinase Inhibition

The enriched mononuclear cell assay, the protocol of which is set forth below, begins with cryopreserved Human Peripheral Blood Mononuclear Cells (HPBMCs) (Clonetics Corp.) that are rinsed and resuspended in a warm mixture of cell growth media. The resuspended cells are then counted and seeded at 1x10⁶ cells/well in a 24-well microtitre plate. The plates are then placed in an incubator for an hour to allow the cells to settle in each well. After the cells have settled, the media is aspirated and new media containing 100 ng/ml of the cytokine stimulatory factor Lipopolysaccharide (LPS) and a test chemical compound is added to each well of the microtiter plate. Thus, each well contains HPBMCs, LPS and a test chemical compound. The cells are then incubated for 2 hours, and the amount of the cytokine Tumor Necrosis Factor Alpha (TNF- α) is measured using an Enzyme Linked Immunoassay (ELISA). One such ELISA for detecting the levels of TNF- α is commercially available from R&D Systems. The amount of TNF- α production by the HPBMCs in each well is then compared to a control well to determine whether the chemical compound acts as an inhibitor of cytokine production.

LPS induced cytokine synthesis in HPBMCs

Cryopreserved HPBMC (cat#CC-2702 Clonetics Corp)

LGM-3 media (cat#CC-3212 Clonetics Corp)

LPS stock 10 g/ml (Cat. No. L 2630 serotype 0111:B4 Sigma)

Human TNF- ELISA (R&D Systems)

DNase I (10mg/ml stock)

Preparation of cells.

LGM-3 media warmed to 37°C.

5 l of DNase I stock added to 10ml media.

Cells thawed rapidly and dispersed into above.

5 Centrifuge 200xg x10min @ RT.

Pellet up in 10ml sterile PBS.

Centrifuge 200xg x10min @ RT.

Pellet resuspended in 10ml LGM-3 then diluted to 50ml with LGM-3.

Perform cell count.

10 Adjust to 1xE06 cells/well.

Seed 1ml/well of a 24 well plate.

Place plate in incubator to plate down for 1 hour.

Preparation of incubation media.

15 LGM-3 containing 100ng/ml LPS (e.g. 50ml media plus 0.5ml LPS stock)

Aliquot into 2ml aliquots and add 1000X inhibitor dilutions.

Incubation

When cells have plated down aspirate media away and overlay with 1ml relevant
20 incubation media. Return plate to incubator for 2 hours or 24 hours. Remove supernatants
after incubation to a labeled tube and either perform TNF (or other) ELISA immediately or
freeze for later assay.

IC₅₀ values are calculated using the concentration of inhibitor that causes a 50% decrease
as compared to a control.

25 Administration and Use

The compounds of the invention are useful among other indications in treating conditions
associated with inflammation. Thus, the compounds of formula (1) or their
pharmaceutically acceptable salts are used in the manufacture of a medicament for
prophylactic or therapeutic treatment of mammals, including humans, in respect of
30 conditions characterized by excessive production of cytokines and/or inappropriate or

unregulated cytokine activity on such cells as cardiomyocytes, cardiofibroblasts and macrophages.

The compounds of the invention inhibit the production of cytokines such as TNF, IL-1, IL-6 and IL-8, cytokines that are important proinflammatory constituents in many different disease states and syndromes. Thus, inhibition of these cytokines has benefit in controlling and mitigating many diseases. The compounds of the invention are shown herein to inhibit a member of the MAP kinase family variously called p38 MAPK (or p38), CSBP, or SAPK-2. The activation of this protein has been shown to accompany exacerbation of the diseases in response to stress caused, for example, by treatment with lipopolysaccharides or cytokines such as TNF and IL-1. Inhibition of p38 activity, therefore, is predictive of the ability of a medicament to provide a beneficial effect in treating diseases such as Alzheimer's, coronary artery disease, congestive heart failure, cardiomyopathy, myocarditis, vasculitis, restenosis, such as occurs following coronary angioplasty, atherosclerosis, IBD, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, multiple sclerosis, acute respiratory distress syndrome (ARDS), asthma, chronic obstructive pulmonary disease (COPD), silicosis, pulmonary sarcosis, sepsis, septic shock, endotoxic shock, Gram-negative sepsis, toxic shock syndrome, heart and brain failure (stroke) that are characterized by ischemia and reperfusion injury, surgical procedures, such as transplantation procedures and graft rejections, cardiopulmonary bypass, coronary artery bypass graft, CNS injuries, including open and closed head trauma, inflammatory eye conditions such as conjunctivitis and uveitis, acute renal failure, glomerulonephritis, inflammatory bowel diseases, such as Crohn's disease or ulcerative colitis, graft vs. host disease, bone resorption diseases like osteoporosis, type II diabetes, pyresis, psoriasis, cachexia, viral diseases such as those caused by HIV, CMV, and Herpes, and cerebral malaria.

Within the last several years, p38 has been shown to comprise a group of MAP kinases designated p38- α , p38- β , p38- γ and p38- δ . Jiang, Y., *et al.*, *J Biol Chem* (1996) 271:17920-17926 reported characterization of p38- β as a 372-amino acid protein closely related to p38- α . In comparing the activity of p38- α with that of p38- β , the authors state that while both are activated by proinflammatory cytokines and environmental stress, p38- β was preferentially activated by MAP kinase kinase-6 (MKK6) and preferentially

activated transcription factor 2, thus suggesting that separate mechanisms for action may be associated with these forms.

Kumar, S., *et al.*, *Biochem Biophys Res Comm* (1997) 235:533-538 and Stein, B., *et al.*, *J Biol Chem* (1997) 272:19509-19517 reported a second isoform of p38- β , p38- β 2, containing 364 amino acids with 73% identity to p38- α . All of these reports show evidence that p38- β is activated by proinflammatory cytokines and environmental stress, although the second reported p38- β isoform, p38- β 2, appears to be preferentially expressed in the CNS, heart and skeletal muscle compared to the more ubiquitous tissue expression of p38- α . Furthermore, activated transcription factor-2 (ATF-2) was observed to be a better substrate for p38- β 2 than for p38- α , thus suggesting that separate mechanisms of action may be associated with these forms. The physiological role of p38- β 1 has been called into question by the latter two reports since it cannot be found in human tissue and does not exhibit appreciable kinase activity with the substrates of p38- α .

The identification of p38- γ was reported by Li, Z., *et al.*, *Biochem Biophys Res Comm* (1996) 228:334-340 and of p38- δ by Wang, X., *et al.*, *J Biol Chem* (1997) 272:23668-23674 and by Kumar, S., *et al.*, *Biochem Biophys Res Comm* (1997) 235:533-538. The data suggest that these two p38 isoforms (γ and δ) represent a unique subset of the MAPK family based on their tissue expression patterns, substrate utilization, response to direct and indirect stimuli, and susceptibility to kinase inhibitors.

Various results with regard to response to drugs targeting the p38 family as between p38- α and either the putative p38- β 1 or p38- β 2 or both were reported by Jiang, Kumar, and Stein cited above as well as by Evers, P.A., *et al.*, *Chem and Biol* (1995) 5:321-328. An additional paper by Wang, Y., *et al.*, *J Biol Chem* (1998) 273:2161-2168 suggests the significance of such differential effects. As pointed out by Wang, a number of stimuli, such as myocardial infarction, hypertension, valvular diseases, viral myocarditis, and dilated cardiomyopathy lead to an increase in cardiac workload and elevated mechanical stress on cardiomyocytes. These are said to lead to an adaptive hypertrophic response which, if not controlled, has decidedly negative consequences. Wang cites previous studies which have shown that in ischemia reperfusion treated hearts, p38 MAPK activities are elevated in association with hypertrophy and programmed cell death. Wang shows in the cited paper that activation of p38- β activity results in hypertrophy, whereas activation of p38- α activity leads to myocyte apoptosis. Thus, selective inhibition of p38- α activity

as compared to p38- β activity will be of benefit in treating conditions associated with cardiac failure. These conditions include congestive heart failure, cardiomyopathy, myocarditis, vasculitis, vascular restenosis, valvular disease, conditions associated with cardiopulmonary bypass, coronary artery bypass, grafts and vascular grafts. Further, to the extent that the α -isoform is toxic in other muscle cell types, α -selective inhibitors would be useful for conditions associated with cachexia attributed to TNF or other conditions such as cancer, infection, or autoimmune disease.

Thus, the invention encompasses the use of compounds which selectively inhibit the activity of the p38- α isoform for treating conditions associated with activation of p38- α , in particular those associated with cardiac hypertrophy, ischemia or other environmental stress such as oxidation injury, hyperosmolarity or other agents or factors that activate p38- α kinase, or cardiac failure, for example, congestive heart failure, cardiomyopathy and myocarditis.

The manner of administration and formulation of the compounds useful in the invention and their related compounds will depend on the nature of the condition, the severity of the condition, the particular subject to be treated, and the judgement of the practitioner; formulation will depend on mode of administration. As the compounds of the invention are small molecules, they are conveniently administered by oral administration by compounding them with suitable pharmaceutical excipients so as to provide tablets, capsules, syrups, and the like. Suitable formulations for oral administration may also include minor components such as buffers, flavoring agents and the like. Typically, the amount of active ingredient in the formulations will be in the range of 5%-95% of the total formulation, but wide variation is permitted depending on the carrier. Suitable carriers include sucrose, pectin, magnesium stearate, lactose, peanut oil, olive oil, water, and the like.

The compounds useful in the invention may also be administered through suppositories or other transmucosal vehicles. Typically, such formulations will include excipients that facilitate the passage of the compound through the mucosa such as pharmaceutically acceptable detergents.

The compounds may also be administered topically, for topical conditions such as psoriasis, or in formulation intended to penetrate the skin. These include lotions, creams, ointments and the like which can be formulated by known methods.

The compounds may also be administered by injection, including intravenous, intramuscular, subcutaneous or intraperitoneal injection. Typical formulations for such use are liquid formulations in isotonic vehicles such as Hank's solution or Ringer's solution.

- 5 Alternative formulations include nasal sprays, liposomal formulations, slow-release formulations, and the like, as are known in the art.


Any suitable formulation may be used. A compendium of art-known formulations is found in Remington's Pharmaceutical Sciences, latest edition, Mack Publishing Company, Easton, PA. Reference to this manual is routine in the art.

- 10 The dosages of the compounds of the invention will depend on a number of factors which will vary from patient to patient. However, it is believed that generally, the daily oral dosage will utilize 0.001-100 mg/kg total body weight, preferably from 0.01-50 mg/kg and more preferably about 0.01 mg/kg-10 mg/kg. The dose regimen will vary, however, depending on the conditions being treated and the judgment of the practitioner.

- 15 It should be noted that the compounds of formula (1) can be administered as individual active ingredients, or as mixtures of several embodiments of this formula. In addition, the inhibitors of p38 kinase can be used as single therapeutic agents or in combination with other therapeutic agents. Drugs that could be usefully combined with these compounds include natural or synthetic corticosteroids, particularly prednisone and its derivatives,
20 monoclonal antibodies targeting cells of the immune system, antibodies or soluble receptors or receptor fusion proteins targeting immune or non-immune cytokines, and small molecule inhibitors of cell division, protein synthesis, or mRNA transcription or translation, or inhibitors of immune cell differentiation or activation.

- As implied above, although the compounds of the invention may be used in humans, they
25 are also available for veterinary use in treating animal subjects.


The following examples are intended to illustrate but not to limit the invention, and to illustrate the use of the above Reaction Schemes.

Example 16-Methoxy-(4-benzylpiperidiny1)-indole-5-carboxamide-3-glyoxalic acid

- 5 0.348 mg (1 mmol) of 6-methoxy-(4-benzylpiperidiny1)-indole-5-carboxamide was dissolved in 15 mL dry dichloromethane and was cooled to 0°C in an ice bath. 0.6 mL of a 2 molar solution of oxalylchloride in dichloromethane (Aldrich) was added dropwise using a syringe under inert atmosphere and the mixture was stirred at 0°C for an h. The ice bath was removed and the mixture stirred further an h. at room temperature. The
- 10 solvent was evaporated and the residue dried under vacuum for 30 Min. The solid obtained was dissolved in a mixture of THF/water and basified with 20 % aq. NaOH. The solvents were removed and the residue dissolved in water and acidified with conc. HCl. The precipitated solid was collected by filtration, dried and recrystallized from ethanol/water to yield 350 mg of the title compound.


15

ESMS. 421, M⁺

Example 26-Methoxy-(4-benzylpiperidinyl)-5-carboxamido-indole-3-glyoxalicacid-4-methylpiperazinamide

5

This compound was prepared using the same procedure used above for the corresponding acid, but substituting 4-methylpiperazine for aq. NaOH and carrying out the reaction in dry dichloromethane instead of THF/water.


10 ESMS. 503, M⁺Example 36-Methoxy-(4-benzylpiperidinyl)-5-carboxamido-indole-3-glyoxalicacid-1-(2-aminoethylpyrrolidine)-amide.

15 This compound was prepared using the same procedure used above, but substituting 1-(2-aminoethyl)-pyrrolidine for 4-methylpiperazine.

MS. M^+ , 517,

Example 4

4-benzylpiperidiny-5-carboxamido-indole-3-glyoxalicamide.



5


0.318g, 1mmol, of 4-benzylpiperidiny-indole-5-carboxamide was dissolved in dry dichloromethane and was reacted with 0.6 mL 2 molar solution of oxalylchloride at 0°C for 1h under nitrogen. Cooling was removed and the mixture was stirred an additional 1h. At RT. The solvent was evaporated and the residue dried under vacuum for 30 Minutes. The product was redissolved in THF and excess of con. Ammonium hydroxide was added. After stirring for 1h. The solvent was removed and the residue recrystallized from ethylacetate-hexane.

10

Yield; 220 mg. MS. M^+ , 389; 345, $M^+ - \text{CONH}_2$

Example 5

6-Chloro-(4'-fluoro-4-benzylpiperidiny)-5-carboxamido-indole-3-glyoxalicacid,
4-methylpiperazinamide.



15

Prepared using similar procedure above described.


MS. M^+ , 524.

5

Example 6

Preparation of 6-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide-3-glyoxalic acid methylester

10



15


This compound was prepared using the same procedure described for the parent glyoxalic acid, but substituting methanol for sodium hydroxide in THF/water.

20 ESMS: M^+ 435.

Example 7

Preparation of 1-methyl-6-methoxy-[4'-fluoro-(4-benzylpiperidinyl)-]-indole-5-carboxamide-3-glyoxalic acid methylester.

28




O.435 g of 6-methoxy- [4'-fluoro-(4-benzylpiperidinyl)-]-indole-5-carboxamide-3-glyoxalic acid methylester was dissolved in 10 mL dry DMF and was cooled to 0°C in an ice-bath. 80 mg NaH (60% dispersion) was added and the mixture stirred for 15 minutes at 0°C and 30 min. at room temperature under inert atmosphere. The reaction mixture was cooled to 0°C and 200 μ L of iodomethane was added. After 30 Min. at 0°C, it was allowed to warm to room temperature and stirring continued for another 4 h. The reaction mixture was poured in to water and extracted with dichloromethane (3 x 50 mL). The extract was dried, evaporated and purified by chromatography on silica gel with ethylacetate-hexane (50 to 90% ethylacetate, gradient).
Yield: 70 %

MS: M⁺, 466.

Example 8

Preparation of 1-methyl-6-methoxy-[4'fluoro-(4-benzylpiperidinyl)-]-indole-5-carboxamide-3-glyoxalic acid.




0.24g (0.51 mmol) of 1-methyl-6-methoxy-[4'fluoro-(4'benzylpiperidiny)]-indole-5-carboxamide-3-glyoxalic acid methylester was dissolved in THF (10 mL) and 1 mL 20 % aq. sodiumhydroxide was added and stirred for 4h. It was diluted with water (5 mL) and stirring continued for 1h. THF was removed under reduced pressure and the remaining solution was diluted with water and acidified with conc. HCl and the product was collected by filtration. It was dried *in vacuo* and recrystallized from ethylacetate.

Yield: 180 mg

ESMS: M⁺, 453.

Example 9


3-(-Methoxybenzoyl)-(4-benzylpiperidiny)-indole-5-carboxamide.



Acylation at 3-position of the indole ring system was achieved by a procedure of C. X.


Yang, *et al.* (Synth. Commun. 27 (12), 2125 (1997). To a solution of 0.318 g (1.0 mmol) of 5(4-benzylpiperidiny)-indole carboxamide in dichloromethane was added 2.2 mL (2.2 mmol) of 1 M solution of ZnCl₂ in ether followed by the dropwise addition of EtMgBr (1.0 mmol). The mixture was stirred for 1h and p-anisoyl chloride (180 mg, 1.1 mmol) was added. The suspension was stirred for 1h and AlCl₃ (0.05 mmol) was added. The resultant mixture was stirred for 2 h and was quenched with Sat. NH₄Cl solution. The organic layer was washed with aq. NaHCO₃ and brine, dried over MgSO₄ and filtered. The solvent was evaporated and the product was purified by Silica gel chromatography.

MS. M⁺, 452.

Example 103-Benzoyl-5-(4-benzylpiperidiny)-indole carboxamide.

Prepared using the same procedure described above, but substituting benzoyl chloride for
5 p-anisoyl chloride.


MS: M^+ , 422.

Example 113-acetyl-5-(4-benzylpiperidiny)-indole-5-carboxamide.

10

Prepared using the same procedure described above, but substituting acetyl chloride for p-anisoyl chloride.

MS: M^+ , 360.

Example 12Preparation of 3-(2-hydroxyacetyl)-5-(4-benzylpiperidiny1)-indole carboxamide

10

3-(2-chloroacetyl)-5-(4-benzylpiperidiny1)-indole carboxamide was prepared using the same procedure described before, but substituting chloroacetyl chloride for p-anisoyl chloride.


Hydrolysis of the chloroacetyl moiety to the hydroxyacetyl was achieved by using published procedure. (J. Org. Chem. 1988, 53, 5446). To a solution of 50 mg (0.13 mmol) of 3-(2-chloroacetyl)-5-(4-benzylpiperidine)-indole-carboxamide in dioxane (3 mL) was added 5 mL of formamide-water (10:1). The reaction mixture was heated at 110°C for 5 h and cooled to RT. The reaction mixture was quenched with sat. NH₄Cl solution, extracted with dichloromethane, dried (MgSO₄) and concentrated. The residue was purified by preparative TLC developed with dichloromethane-methanol (20:1) to give 20 mg (42%) of the title compound.

20

MS. M⁺, 376.

Example 135-(4-Benzylpiperidiny1)-carboxyamido-indole-3-(3'-oxo)-ethylpropionate.

25



The acylation of 3-position was obtained by published method. [Synth. Commun. 1977, 27
(12), 2125]. To a solution of 3.6 mL of ZnCl_2 (3.6 mmol, 1M solution in diethyl ether) in
5 THF (15 mL) was added n-BuLi (2.2 mL, 3.6 mmol) dropwise at 0°C . A white
suspension was formed during the addition. The reaction mixture was warmed to RT,
stirred for 1h, and a solution of 5-(4-benzylpiperidinyI)-indolecarboxamide in
dichloromethane (10 mL) was added. The resulting mixture was stirred for 1h and ethyl 3-
10 chloro-3-oxopropionate (585 mg, 3.9 mmol) was added. After 1h, the reaction mixture was
quenched with saturated NH_4Cl , extracted with dichloromethane, dried (MgSO_4) and
concentrated. The residue was purified by silica gel chromatography to yield 200 mg of
the desired product.


15 MS> M^+ , 431.

Compounds 1-55 of Figure 2 were similarly prepared.

Example 14

Synthesis of (2S,5R)N-4-Fluorobenzyl-*trans*-2,5-dimethylpiperazine


(+/-)N-Benzyl-*trans*-2,5-dimethylpiperazine was synthesized as follows




50 g *trans*-2,5 dimethylpiperazine was dissolved in 300 mL ethanol and treated with 26.36
mL (1/2 equivalents) benzylbromide. The mixture was stirred at room temperature for 12
25 hours and concentrated. The residue was taken up in ethyl acetate and washed with 10%
aqueous sodium bicarbonate and saturated sodium chloride; dried over anhydrous

magnesium sulfate and concentrated to give crude 1-benzyl-*trans*-2,5-dimethylpiperazine, as an oil.

This material was chromatographed using DCM/MeOH 95/5 to remove the di-alkylated product and then with DCM/MeOH/TEA 90/10/0.1 to elute the benzyl-*trans*-2,5-dimethylpiperazine. 19.6 g of the pure product was obtained as an oil.



To a solution of (+/-)-N-benzyl-*trans*-2,5-dimethylpiperazine (59 g, 0.29 mol) in Methanol (150 mL) was added a solution of (+) tartaric acid (87 g, 0.58 mol) in Methanol (250 mL) dropwise over 5 min. Crystallization is effected by keeping the resulting mixture at 0 °C for 48-72 hours. Scratching of the solution after 12-16 hours facilitates the crystallization process. The mixture was filtered and washed with cold Methanol and dried to give the ditartaric acid salt (73.9 g) as white crystals. A single recrystallization from Methanol, cooling to room temperature afforded the salt as white crystals. (58 g) $[\alpha]_D = +47$, ($c = 1.00$, Methanol).



31 g, 15.27 mmol, dimethylbenzylpiperazine was treated with 43 g, 19.85 mmol of di-tert-butyl-dicarbonate in 250 mL THF for 4 hours. The reaction was monitored by TLC and is


essentially complete when the addition of (BOC)₂O is complete. The solvent was removed and the residue was taken up in ethyl acetate and washed with 10% aqueous sodium carbonate and saturated sodium chloride, dried over anhydrous sodium sulfate and concentrated to give 39.3 g of the Boc-protected compound. This material was used for the next step without further purification.

39.3 g, 131 mmol, of the Boc-protected benzylpiperazine was treated with 3.93 g of Pearlman's catalyst, in 150 mL methanol with 3 mL acetic acid for 4 hours at 40 psi, hydrogen pressure in a parr shaker. The reaction mixture was filtered through celite and concentrated to give a residue that was dried under high vacuum and then dissolved in


250 mL dry ethanol and treated with 1.2 equivalents, 158 mmol of 4-Fluorobenzylbromide, and 2 equivalents, 262 mmol of triethylamine for 5 hours. The reaction mixture was monitored by TLC and found to be complete at that time. The solvent was removed and the residue was taken up in ethyl acetate and washed with 10% aqueous sodium carbonate and saturated sodium chloride, dried over anhydrous sodium sulfate and concentrated to give 47 g of crude product. This crude material was chromatographed on silica gel using Hexane/Ethyl acetate 95/5 to elute the product. After chromatography 29 g of the 4-Fluorobenzyl-trams-2,5-dimethylpiperazine was obtained.

20 g of the 1-(4-Fluoro)benzyl-trams-2,5-dimethylpiperazine, 62 mmol was treated with a mixture 200 mL 4 N HCl/Dioxane/2 M HCl/Ether (1:3) for 1 hour. Electron impact mass spectroscopy confirmed the formation of product and the disappearance of starting material at that time. The reaction mixture was concentrated to give a white solid. This material was repeatedly washed with hexane and ether to remove residual dioxane, and then dried under vacuum extensively before use. 17.6 g of the final product was obtained as a white fluffy solid.

35

Example 15Synthesis of Chiral Compounds

5



The 5-carboxylic-6-chloro indole acid (1.56 g, 7.44 mmol) was dissolved in dry methylene chloride 60 mL to this was added the EDAC.HCl (1.57 g, 8.18 mmol) and DMAP(10 % mol). After stirring under nitrogen for 10 min. the amine (2.19 g, 7.5 mmol) was added, followed by triethylamine (3 mL, 21.52 mmol). After overnight at room temperature, the reaction mixture was concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate, saturated sodium chloride, dried over anhydrous sodium sulfate and filtered. Concentration gives the crude product that was chromatographed on silica gel using a gradient of EtOAc/Hexane 2/8 – 6/4. TLC R_f 0.435 (EtOAc:Hexane, 1:1), EIMS M^+ 413.


1.02 g of the product from the above step was dissolved in 30 mL dry DCM. The reaction mixture was purged with nitrogen and placed in an ice bath. To this was added 4 mL of 2M oxalyl chloride in DCM. The reaction mixture was stirred at room temperature for 1 hour and then at room temperature for 2 hours. Reaction mixture was concentrated on a rotary evaporator. After drying on a vacuum pump for 15 min. the residue (a yellow solid) was dissolved in dry DCM, 30 mL, to which was added 4 mL of a 2M solution of dimethylamine in THF. 30 minutes later the reaction mixture was concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate,

saturated sodium chloride, dried over anhydrous sodium sulfate and filtered.

Concentration gives the crude product that was chromatographed on silica gel using a gradient of EtOAc 100% - EtOAc/MeOH 9:1. TLC R_f 0.5 (EtOAc:MeOH, 9:1), EIMS M^+ 513.

- 5 The white solid from the above step was dissolved in 10 mL dry DCM. To this was added sufficient 2 M HCl in ether, till a precipitate persists. The mixture was then concentrated on a rotary evaporator to dryness and then dried overnight under high vacuum to give the final product (1.08 g).

ADDITIONAL EXAMPLES



10

- Synthesis of 2:** Methyl 4-amino-2-chlorobenzoate (1) (18.5g) was dissolved in dichloromethane (350ml) and methyl thioacetaldehyde dimethylacetal (13.6g) was added. The mixture was cooled to -45°C (dry ice/acetonitrile bath). N-chlorosuccinimide (16.0g) in 350ml dichloromethane was added dropwise over 1 hr 30min while maintaining bath temp at -45°C. The reaction mixture was stirred additional 1 hr, then triethylamine (16mL, 100 mmols) in 30ml dichloromethane was added dropwise over 5 min, reaction was warmed to room temp, then refluxed for 16h. Solvent was removed and residue taken up in 500ml carbon tetrachloride, triethylamine-hydrochloric acid was removed by filtration, filtrate was heated to reflux for 2 h. Solvent was removed by rotary evaporation.

- The residue was dissolved in 250ml tetrahydrofuran and 250ml 10% hydrochloric acid was added. The mixture was stirred overnight at room temperature until the complete disappearance of the starting material was observed. Solvent was removed under vacuum, acidic aqueous solution was extracted with ethyl acetate (3 x 125ml). The combined ethyl

25

acetate extracts were washed with 10% hydrochloric acid, water and dried over anhydrous sodium sulfate. Solvent was removed under vacuum. Crude product mixture was purified on a silica column eluting with ethyl acetate:hexanes (15:85) to give 6.4g of the desired product 2.


5

Synthesis of 3: Methyl 6-Chloro-3-thiomethyl-5-indole carboxylate (5.2g) was dissolved in 150ml ethanol:tetrahydrofuran (9:3) and treated with Raney-Nickel. Reaction was monitored by mass spec at 30min intervals, with subsequent addition of Raney-Nickel until reaction was complete. When reaction was complete reaction was carefully filtered through celite and the celite washed with methanol several times and filtrate evaporated. Residue was taken up in ethyl acetate, washed with water, dried over anhydrous sodium sulfate. The solvent was removed to give 3 (3.2g).

10

Synthesis of 4: Methyl ester 1.5g was dissolved in 30ml methanol/water 50:50. The reaction mixture was heated at 50°C for 2 h with 4 Mol. equivalent sodium hydroxide. The reaction mixture was cooled in ice-bath, acidified to pH 3 with 5M hydrochloric acid,. Removed methanol by rotary evaporation and extracted with ethyl acetate. The extract was washed with saturated sodium chloride and dried over anhydrous sodium sulfate. Evaporation of the solvent gave the desired acid 4 (1.48 g).

20



Synthesis of 5:

STEPA: The phosphonate A (38.4 g) and the piperidone B (35.4) were dissolved in anhydrous dimethylformamide (400 mL). To this sodium hydride (60% suspension in oil) was added in portions while the reaction is maintained at 0°C. After the addition of sodium hydride was complete the reaction mixture was stirred for 30 min. and then the ice bath was removed, the reaction was allowed to stir for 6h as it slowly warmed to ambient temperature. The reaction was again cooled in an ice bath and quenched with methanol.

Water was added to the reaction mixture, and the product extracted with ethyl acetate. The


30

ethyl acetate layer was washed with saturated sodium chloride and dried over anhydrous magnesium sulfate. The solvent was removed to give the crude alkene, which is purified by column chromatography eluting with ethyl acetate/hexane (1:9) to give 21.8 g of the desired product C.

5

STEP B: 10.1 g of C was dissolved in 50 mL methanol. After purging the solution with nitrogen, 5% Palladium on carbon (1g) catalyst was added followed by 1 mL acetic acid. The reaction mixture was hydrogenated for 4 h at 40-50 psi. The reaction mixture was filtered through celite and concentrated. The residue was treated with 2 M hydrochloric acid in ether to convert to the hydrochloric acid salt. The white solid that was obtained was dried under vacuum, extensively, to give 7.8 g of 5 as the hydrochloric acid salt.

10



15

Synthesis of 6: A mixture of 6-chloro-indole-5-carboxylic acid (1.95 g), 4-fluorobenzylpiperidine hydrochloric acid salt (2.76g) was taken in 50 mL dry dichloromethane and was treated with triethylamine (1.7 mL). The mixture was stirred until a clear solution was obtained. 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (2.3g) and dimethylaminopyridine (0.25 g) was added and the mixture was stirred for 20h at ambient temperature. The mixture was poured in to water and the organic layer separated. The aq. Solution was further extracted twice with dichloromethane. The combined extract was washed with 10% hydrochloric acid. The organic layer was dried over magnesium sulfate


20

and evaporated. The product was purified by chromatography on silica gel eluting with ethylacetate:hexane (3:7) to give 6 (2.78 g).

Synthesis of 7: 8.61 g of 6-chloro-(4-F-benzylpiperidinyl)- indole-5-carboxamide (6) was dissolved in about 100 mL dry DMF and the solution was cooled in an ice-bath. To the cold solution was added 30 mL of 1M solution of sodium bis-(trimethylsilyl)-amide in tetrahydrofuran under inert atmosphere. The reaction was allowed to stir at 0°C for 15 min and at ambient temperature for another 30 min. The reaction mixture was cooled again in ice-bath and 2.5 mL of iodomethane was added. After stirring for 30 min. at 0°C, it was allowed to warm to AMBIENT TEMPERATURE and stirring continued for 18h at AMBIENT TEMPERATURE. The mixture was diluted with water (or brine) and the product was extracted with ethylacetate (4 x 75 mL). The combined extract was washed with water and dried over anh. MgSO₄ . The solvent was removed and the product was purified by column chromatography on silica gel eluting with ethylacetate:hexane (1:4) to yield 8.0 g of the desired product 7.

Synthesis of 8: 8g of 7 was dissolved in about 100 mL of anhydrous dichloromethane and was cooled in an ice-bath. To this a 2M solution of oxalylchloride (20.8 mL) in dichloromethane was added slowly via a syringe and the mixture was allowed to stir at 0°C for 1h. The ice-bath was removed and stirring continued for an additional 2h at AMBIENT TEMPERATURE. The solvent was removed under reduced pressure and the residue was pumped for 15 min. to remove any excess oxalylchloride present. The product was immediately redissolved in anh. dichloromethane (150 mL), cooled in an ice-bath, 30 mL of a 2M solution of dimethylamine in tetrahydrofuran was added rapidly via a syringe. After 15 min. stirring it was allowed to stir for another 15 min. at ambient temperature. The solution was washed with water to remove the salt, and dried. After evaporation, the residue was purified on silica gel, eluting with chloroform:methanol (99:1) to yield 9.3g of 8.

40

**Synthesis of 10:**


Synthesis of 10:

STEP A: To a solution of dimethyl piperazine **9** (25g) in 300 ml of absolute ethanol was added 400 ml of 2N hydrogen chloride in diethyl ether. The solution was warmed to 70 °C in an oil bath for 20 minutes. The solution was then cooled to room temperature and set at 6 °C overnight. The solid obtained, was collected by filtration. Yield 39.8 g (dihydrochloride salt of trans-2,5 dimethylpiperazine) after drying overnight under high vacuum.

STEP B: An ethanol solution of 42.9g of dimethyl piperazine dihydrochloride the from STEP A and 26.1g *trans*-2,5 dimethylpiperazine was vigorously stirred in an oil bath at 80 °C until all starting materials were dissolved. The temperature of oil bath was reduced to 65 °C and 33.1g of 4-fluoro benzylchloride was added. After stirring at this temperature for 30 min., the solution was placed in a 6 °C refrigerator overnight. The solid was removed from the solution by filtration and excess of 2N hydrogen chloride in diethyl ether was added to the filtrate. The filtrate was kept at 6 °C overnight and the solid collected. The solid was suspended in 5% sodium hydroxide aqueous solution and extracted three times with ethyl acetate. The organic layer was dried over sodium sulfate and dried down to give a yellow oil.

STEP C: A solution of 50.7 g (L)-tartaric acid in 130 ml of boiling methanol was added to 70 ml of hot methanol solution of 37.5 g of the product from STEP B. The solution was set at 6 °C for 96 hours before collection of white fine crystals by filtration. This material was recrystallized from boiling methanol. The product was collected by filtration after being kept at a 6 °C overnight. Yield 30.5 g of ditartaric acid salt ($[\alpha] = +43.2^\circ$, $c=1$).

41




Synthesis of 11: The indole ester 3 (0.526 g) was dissolved in 10 mL acetone (dry) and placed in an ice bath. To this was added crushed potassium hydroxide (0.7 g, 12.5 mmol), after stirring for five minutes at 0°C, methyl iodide (400 μ L, 6.272 mmol) was added to the reaction mixture. The reaction mixture was stirred at 0 °C for 10 minutes and then at room temperature for 30 minutes. After removal of the solvent, the residue was taken up in ethyl acetate and washed with saturated sodium chloride. After drying over anhydrous sodium sulfate, filtration and rotary evaporation, a solid was obtained, 0.7 g. Column chromatography on silica with ethyl acetate:hexane (2:8) gave methyl ester of 11 as a white solid. 0.52 g. 0.52 g of this product was dissolved in 50 mL methanol and treated with 5 mL 10 N sodium hydroxide, the reaction mixture was heated at 50°C for 2 h. The reaction mixture was cooled to room temperature and concentrated to a solid on a rotary evaporator. The residue was taken in 50 mL water, washed with ether and placed in an ice bath. The basic solution was acidified with 10% hydrochloric acid to pH 2. The precipitate was extracted with ethyl acetate and the ethyl acetate layer was washed with saturated sodium chloride solution. Drying over anhydrous sodium sulfate, filtration and concentration on a rotary evaporator to gave 11 (0.48 g) as a white solid.

Synthesis of 12: 1.56 g of the acid (11) was dissolved in dry methylene chloride 10 mL and to this was added the 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (1.57 g) and dimethylaminopyridine (catalytic). After stirring under nitrogen for 10 min., 2.19 g of the amine (10) was added, followed by triethylamine (3mL). After stirring overnight at room

temperature, the reaction mixture was concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate, aq. saturated sodium chloride, dried over anhydrous sodium sulfate and filtered. Concentration gave the crude product which was chromatographed on silica gel using ethyl acetate:hexane (4:6) as eluent to give 1.02g of the desired product.

Synthesis of 13: 1.02 g of 12 from the previous step was dissolved in 30 mL dry dichloromethane. The reaction mixture was purged with nitrogen and placed in an ice bath. To this was added 4 mL of 2M oxalyl chloride in dichloromethane. The reaction mixture was stirred at 0°C for 1 hour and then at room temperature for 2 h. Reaction mixture was concentrated on a rotary evaporator. After drying on a vacuum pump for 15 min. the residue (a yellow solid) was dissolved in dry dichloromethane (30 mL) to which was added 4 mL of a 2M solution of dimethylamine in tetrahydrofuran. 30 minutes later the reaction mixture was concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate, saturated sodium chloride, dried over anhydrous sodium sulfate and filtered. Concentration gives the crude product that was chromatographed on silica gel with ethyl acetate:methanol (9:1). The white solid obtained was dissolved in 10 mL dry dichloromethane. To this was added sufficient 2 M hydrochloric acid in ether, till a precipitate persisted. The mixture was then concentrated on a rotary evaporator to dryness and then further dried overnight under high vacuum to give 13 (1.08 g).



Synthesis of 15: To a solution of aniline **1** (9.25 g, 0.05 mol) and pyruvic aldehyde dimethyl acetal **14** (11.8 g, 0.1 mol) in 200 mL glacial acetic acid was added anhydrous sodium sulfate (71.0g, 0.5 mol) and the mixture was stirred for 30 min. Powdered sodium triacetoxy borohydride (31.8 g, 0.15 mol) was then added in portions for a period of 5 min.


5 The reaction mixture was stirred for an additional 2 h. Acetic acid was removed under reduced pressure and the residue was made basic by adding sufficient amount of saturated sodium bicarbonate solution. The product was then extracted with ethyl acetate, dried with sodium sulfate and evaporated to get an oil. This was chromatographed on silica gel column using ethyl acetate:hexane (3:7) to give **15** (14 g) as colorless oil.

10 **Synthesis of 16 and 17:** To a suspension of fresh aluminum chloride (18.5 g) in 200 ml dry chloroform at 0°C was added a solution of ketal **15** (13.3 g) in 100 ml chloroform slowly and the mixture was allowed to warm up to the room temperature and stirred overnight. Ice-cold water was added carefully to quench the aluminum chloride and the
15 organic layer was separated and washed with sodium bicarbonate solution, dried and evaporated to get a white solid. The isomers were separated using silica gel column chromatography using ethyl acetate:hexane (1:9). The 6-cholo indole **17** (2.0 g) eluted first followed by 4-chloro isomer **16** (3.8 g).

20 **Synthesis of 18:** To a solution of 1.3 g of indole **17** in 15 mL of methanol was added a solution of 0.9 g of sodium hydroxide in 20 mL of water. The reaction mixture was heated at 50°C for 4 h where upon a clear solution resulted. Cooled and evaporated off methanol and the residue was diluted with water and acidified with 10% hydrochloric acid. The product was extracted with ethyl acetate. The organic layer was dried over sodium sulfate,
25 filtered and evaporated to obtain indole acid **18** (1.2 g) as white solid.

2-Methyl-6-methoxyindole-5-carboxylic acid was also synthesized using the above synthetic procedure.

44




Synthesis of 21: To a solution of methyl 4-amino-6-methoxy 5-benzoate (**19**) (6.0 g, 0.033 mol) and dimethyl acetal **20** (7.0 g, 0.066 mol) in 150 mL glacial acetic acid was added anhydrous sodium sulfate (47.0 g, 0.33 mol) and the mixture was stirred for 30 min. Powdered sodium triacetoxo borohydride (20.1 g, 0.099 mol) was then added in portions for a period of 5 min. The reaction mixture was stirred for an additional 2 h. Acetic acid was removed under reduced pressure and the residue was made basic by adding sufficient amount of saturated sodium bicarbonate solution. The product was then extracted with ethyl acetate, washed with saturated sodium chloride, dried over sodium sulfate and evaporated to get an oil. This was chromatographed on silica gel column using ethyl acetate:hexane (3:7) as eluent and the desired product **21** was obtained (5.2 g) as an oil.

Synthesis of 22: To a solution of **21** (3.6 g) and iodomethane (5.7 g) in 50 mL anhydrous dimethylformamide was added potassium t-butoxide (1.0 M in tetrahydrofuran, 20 mL) at ambient temperature. The reaction mixture was stirred at ambient temperature for 0.5 h and poured into 250 mL ethyl acetate, washed with water (4x100 mL), brine (50 mL) and dried over magnesium sulfate. Evaporation of solvent afforded 3.26 g of **22**. The product was used for next step without purification.

Synthesis of 23: To a suspension of anhydrous aluminum chloride (0.71 g) in 20 mL anhydrous 1,2-dichloroethane was added, dropwise a solution of **22** (1 g) in 10 mL 1,2-dichloroethane with stirring. The reaction was heated to 80°C for 0.5 h. At the end of this time, the reaction mixture was quenched with methanol, solvents evaporated, then ethyl acetate (100 mL) was added. The organic phase was washed with water, aq. sodium

45

bicarbonate and brine and concentrated. The crude product was purified by silica chromatography using ethyl acetate:hexane (3:7) to give **23** 0.22 g.




5

Synthesis of 24. To a suspension of 1.2 g of indole acid **18** and 1.6 g of **5** in 30 mL dichloromethane was added 0.7 g of triethylamine followed by 1.4 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide. The clear solution obtained was stirred for 4 h. The solvent was evaporated and the residue was taken up in ethyl acetate and washed with water, dil. hydrochloric acid, and brine. The organic layer was then dried with sodium sulfate and evaporated. The product was isolated (1.3 g) as white solids after silica gel chromatography using ethyl acetate:hexane (1:4).


Synthesis of 25: A solution of 1.3 g of **24** in 20 mL dichloromethane was cooled to 0°C and a solution of 2 M oxalyl chloride in dichloromethane (3.4 mL, 6.8 mmol) was added and the reaction mixture was stirred at 0°C for 1 h. The temperature was allowed to come to room temperature and continued stirring for an additional 1 h. The solvent was evaporated and the residue was dried under vacuum. The acid chloride was dissolved in dichloromethane (25 mL) and a solution of 6.8 mL (13.6 mmol) of 2 M N,N-dimethylamine in tetrahydrofuran was added all at once. The solvent was removed and the product was purified by silica gel column chromatography using methanol:chloroform (2:98). The product was obtained as white solid (1.4 g).

46



Synthesis of 27: A solution of 26 (130 mg) in anhydrous dimethylfor (5 ml) was stirred at 0 °C under nitrogen. Sodium hydride(25 mg, 60% dispersion in oil) was added and stirred for 5 min., then room temp for 30 min. The mixture was cooled at 0 °C and chloromethyl methyl sulfide (70 μ l) was added. The reaction mixture was stirred for 20 h at room temp, and then was extracted with ethyl acetate. The organic layer was dried over sodium sulfate, concentrated under vacuum. The residue was purified by chromatotron using methanol:chloroform (2:98) to give 100 mg of 27.


Synthesis of 28: A solution of oxone (120 mg) in water (1 ml) was added to the suspension of 27 (100 mg) in glacial acetic acid (4 ml). The reaction mixture was stirred at room temp. for 18 h, washed with water, and concentrated under vacuum. The residue was purified by chromatotron using methanol:chloroform (2:98) to give 60 mg of 28.



Synthesis of 30: To a solution of 29 (200 mg) in tetrahydrofuran (20 mL) was added sodium bis(trimethylsilyl)amide (0.51 mL, 0.51 mmol) dropwise at 0 °C. The mixture became light yellow solution, warmed to ambient temperature slowly and was stirred for 30 min. Then methoxy methyl chloride was added and the reaction mixture was stirred


overnight. The reaction mixture was quenched with ammonium chloride and extracted with dichloromethane. The combined organic layer was washed with brine, dried and concentrated. The residue was purified by chromatography on silica gel and eluted with methanol:dichloromethane (5:95) to give 190 mg of **30**.

5



Synthesis of 31: To a solution of **29** (200 mg) in tetrahydrofuran (10 mL) was added sodium bis(trimethylsilyl)amide (0.56 mL, 0.56 mmol) dropwise at 0 °C. The reaction mixture was warmed to ambient temperature slowly, stirred for 30 min, and then added to the solution of tosyl cyanide (115 mg) in tetrahydrofuran (10 mL). The resulting mixture was stirred for 2 h at rt, quenched with ammonium chloride and extracted with dichloromethane. The combined organic layer was washed with brine, dried and concentrated. The residue was purified by chromatography on silica gel and eluted with methanol:dichloromethane (5:95) to give 90 mg of **31**.


15



Synthesis of 34: Piperidone **33** was added to a solution of 4-fluorobenzyl magnesium chloride (**32**) (33 mL, 8.3 mmol, 0.25 M in ethyl ether) slowly at 0 °C. The reaction mixture was warmed up to ambient temperature and then at reflux for 6 h. The resulting milky solution was treated with ammonium chloride (saturated) and extracted with ether. The combined organic layer was washed with brine, dried and concentrated. The residue was purified by chromatography on silica gel eluting with hexane:ethyl acetate (4:1) to give 990 mg of **34**.

25

Synthesis of 35: The alcohol **34** (670 mg) in dichloromethane (10 mL) was added to the solution of diethylaminosulfur trifluoride (0.57 mL, 4.34 mmol) in dichloromethane (20 mL) at -78 °C. The reaction mixture was warmed up to ambient temperature slowly and stirred for 2 h, and then treated with sodium carbonate (saturated) and extracted with dichloromethane. The combined organic layer was washed with brine, dried and concentrated. The residue was purified by chromatography on silica gel eluting with hexane:ethyl acetate (8:1) to give 300 mg of **35**.



Synthesis of 36: A mixture of the **35** (456 mg) and 4 M of hydrochloric acid in dioxane (10 mL) was stirred for 4 h. The reaction mixture was neutralized with sodium carbonate, extracted with ethyl acetate. The combined organic layer was washed with brine, dried and concentrated. The crude product was used in the next reaction without further purification.

Synthesis of 38: To the suspension of indole carboxylic acid **37** (475 mg) in anhydrous dichloromethane (20 mL) was added piperidine **36** (350 mg, 1.66 mmol). The mixture was stirred for 10 min and then 1-[3-(dimethylaminopropyl)]3-ethylcarbodiimide hydrochloride (475 mg) and dimethylaminopyridine (202 mg) were added. The reaction mixture became clear and was continually stirred for overnight. Then the reaction mixture

was treated with 10% of hydrochloric acid solution, and extracted with dichloromethane. The combined organic extracts were washed with sodium bicarbonate, brine, dried and concentrated. The residue was purified by chromatography on silica gel eluting with dichloromethane:ethyl acetate (9:1) to give 300 mg of **38** as a white foam.

5

Synthesis of 39: To the suspension of **38** (200 mg) was added oxaly chloride (0.52 mL, 1.04 mmol, 2 M in dichloromethane) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 30 min, and then warmed up to rt. and stirred for 5 h. The yellow suspension was formed. The solvent and excess oxaly chloride were removed under the reduced pressure.


10 The yellow solid formed was dried under vacuum, and dissolved in dichloromethane then cooled in ice bath. Dimethyl amine (1.04 mL, 2.08 mmol, 2 M in tetrahydrofuran) was added. After 30 min, the reaction mixture was treated with water and extracted with dichloromethane. The organic extracts were washed with water, brine, dried and concentrated. The residue was purified by chromatography on silica gel eluting with 3%
15 of MeOH in dichloromethane to give 190 mg of **39** as a white solid.

Synthesis of 40: To a solution of **39** (75 mg) in tetrahydrofuran (10 mL) was added KHMDS (0.46 mL, 0.23 mmol, 0.5 M in toluene) dropwise at 0 °C. The mixture was warmed to rt. slowly and stirred for 30 min. Then methyl iodide (33 mg, 0.23 mmol) was
20 added slowly to the reaction mixture, then stirred for 2 h. The reaction mixture was quenched with ammonium chloride and extracted with dichloromethane. The combined organic extracts were washed with brine, dried and concentrated. The residue was purified by chromatography on silica gel eluting with methanol:dichloromethane (2:98) to give 50 mg of **40**.

25

Synthesis of 41: To a solution of **40** (75 mg) in tetrahydrofuran (10 mL) was added KHMDS (0.46 mL, 0.23 mmol, 0.5 M in toluene) dropwise at 0 °C. The mixture was warmed to rt. slowly and stirred for 30 min. Then methoxy methyl chloride (18 mg, 0.23 mmol) was added slowly to the reaction mixture, then stirred overnight. The reaction
30 mixture was quenched with ammonium chloride and extracted with dichloromethane. The combined organic extracts were washed with brine, dried and concentrated. The residue

was purified by chromatography on silica gel eluting with methanol:dichloromethane (2:98) to give 65 mg of **41**.




- 5 **Synthesis of 43:** To a suspension of 6-methoxy-5-indolecarboxylic acid (**37**) (200 mg) in dichloromethane (5 mL) was added 1-[3-(dimethylaminopropyl)]3-ethylcarbodiimide hydrochloride (258 mg, 1.35 mmol) in one portion. The reaction mixture was stirred till all of the solid was dissolved. 4-Benzyl-4-hydroxypiperidine (**42**) (258 mg), obtained by removal of the protecting group from **34** as outlined in the synthesis of **36**, was added and
- 10 stirred at ambient temperature. The reaction mixture became cloudy. A catalytic amount of dimethylaminopyridine (10 mg) was added and stirred at ambient temperature overnight. The reaction mixture was treated with water, and extracted with dichloromethane. The combined organic layer was washed with 10% of hydrochloric acid solution, sodium bicarbonate (saturated) and brine, then dried and concentrated to give 300
- 15 mg (82%) of **43** as a yellow foam.

Synthesis of 44: Oxalyl chloride (0.41 mL, 1.65 mmol, 2 M in dichloromethane) was added to a solution of 6-methoxy-(4-benzyl-4-hydroxypiperidinyl)-indole-5-carboxamide (300 mg) in dichloromethane (20 mL) at 0 °C dropwise. The reaction mixture was stirred

20 at 0 °C for 30 min, and then warmed up to ambient temperature slowly. After 2 h., a yellow precipitation was formed. The reaction mixture was concentrated under the reduced pressure and dried for 1 h under reduced pressure. The resulting yellow solid was suspended in dichloromethane (20 mL). Methyl piperazine (0.2 mL) and diisopropyl ethyl

amine (0.2 mL) were added at rt. The mixture was stirred for 1 h, treated with water, and extracted with dichloromethane. The residue was purified by chromatography on silica gel eluting with dichloromethane:methanol (10:1) to give 180 mg of a white solid. 20 mg of this white solid was dissolved in methanol (1 mL). To this a saturated hydrochloric acid solution in methanol was added dropwise till the pH value was around 3. Then the solvents were removed and the product was dried to give **44**.

Synthesis of 45: To a solution of **44** (65 mg) in anhydrous tetrahydrofuran (10 mL) was added sodium bis(trimethylsilyl)amide (0.25 mL, 0.25 mmol, 1.0 M solution in tetrahydrofuran) dropwise at ambient temperature. The mixture was stirred at ambient temperature for 30 min, and then ethyl chloroformate (0.024 mL, 0.25 mmol) was added to the reaction mixture. After 1 h, the reaction mixture was quenched with ammonium chloride (saturated), and extracted with ethyl acetate. The combined organic layer was washed with brine, dried (magnesium sulfate) and concentrated under reduced pressure. The residue was purified by chromatography on silica gel, eluting with dichloromethane:methanol (10:1) to give 15 mg of the title compound **45** as a white solid. This was dissolved in methanol (1 mL) and to this was added saturated hydrochloric acid solution in methanol dropwise till the pH value is around 3. Then the solvent was removed and the product was dried to give the hydrochloric acid salt of **45**.



Synthesis of 47:

STEPA: Piperizinone **46** (5.0 g) was dissolved in 15 mL tetrahydrofuran, to this was added a solution of di-tertbuty-dicarbonate (12.04 g in 50 mL tetrahydrofuran). After stirring at

room temperature for 4 h. The reaction mixture was concentrated to dryness on a rotary evaporator and the solid obtained used for the next reaction without further purification.


STEP B: 10 g of product from STEP A was dissolved in anhydrous acetone and the reaction mixture was cooled in an ice bath. To this was added crushed potassium
5 hydroxide (14.02 g). After stirring at 0 °C for 10 min. 4-fluorobenzylbromide (23.5 g) was added and the reaction mixture stirred at 0 °C for 20 min and at room temperature for 1 h. After removal of the solvent on a rotary evaporator the residue was taken up in ethyl acetate and washed with 10% aqueous sodium carbonate, brine and dried over anhydrous sodium sulfate. Filtration and concentration gave crude material that was purified by
10 chromatography on silica gel using ethyl acetate/hexanes gave 14 g of **47**.

Synthesis of 48: 14 g of **47** was dissolved in 100 mL methylene chloride and treated with 100 mL 2 M Hydrochloric acid in ether for 2 h. The solvent was removed and the solid obtained was washed with ether and hexanes and dried under high vacuum to give **48**
15 (11.3g) as the hydrochloride salt.

Synthesis of 49: 1g of **48**, was dissolved in 20 mL methylene chloride and treated with 6-methoxy-5-indole carboxylic acid (**37**) (0.87 g), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (0.95 g), triethylamine (700 µL) and
20 dimethylaminopyridine (catalytic) and the reaction mixture stirred overnight. The reaction mixture was concentrated and the residue taken up in ethyl acetate, washed with 10% aqueous sodium carbonate, brine and dried over anhydrous sodium sulfate. Filtration and concentration gave crude material that was purified by chromatography on silica gel using ethyl acetate/hexanes to give **49** (1.6 g).

Synthesis of 50: 1.6 g of **49** was dissolved in 25 mL dry dichloromethane. The reaction mixture was purged with nitrogen and placed in an ice bath. To this was added, 5 mL 2M oxalyl chloride in dichloromethane. The reaction mixture was stirred at room temperature for 1 h and then at room temperature for 2 h. Reaction mixture was concentrated on a
30 rotary evaporator and after drying on a vacuum pump for 15 min. the residue (a yellow solid) was dissolved in dry dichloromethane. To this residue a 2M solution of

dimethylamine in THF (5 mL) was added and the reaction mixture stirred for 0.5h. The reaction mixture was then concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate, saturated sodium chloride, dried over anhydrous sodium sulfate and filtered. Concentration gives the crude product that was chromatographed on silica gel using ethyl acetate/hexanes to give **50** (1.45 g).



Synthesis of 52: 2R, 5S-transdimethyl piperazine **51** (5.0 g) was dissolved in 15 mL ethanol, to this was added 2-bromoethylbenzene (4.4 g). The reaction mixture was stirred at 40 °C for 30 min., cooled to room temperature and concentrated to dryness on a rotary evaporator. The residue was taken up in ethyl acetate and washed with 10% aqueous sodium carbonate, brine and dried over anhydrous sodium sulfate. Filtration and concentration gave crude material that was purified by chromatography on silica gel using ethyl acetate/hexanes to give **52** (5.4 g).


Synthesis of 53: 5 g of **52** was dissolved in 60 mL methylene chloride and treated with 60 mL 2 M hydrochloric acid in ether for 3 h. The solvent was removed and the solid obtained was washed with ether and hexanes and dried under high vacuum to give **53** as the hydrochloride salt, 3.6 g.

Synthesis of 54: **53** (1.32 g), was dissolved in 20 mL methylene chloride and treated with 6-methoxy-5-indole carboxylic acid (**37**) (0.9 g) in the presence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (1.04 g), 800 µL triethylamine and catalytic DMAP,

overnight. The reaction mixture was concentrated and the residue taken up in ethyl acetate, washed with 10% aqueous sodium carbonate, brine and dried over anhydrous sodium sulfate. Filtration and concentration gave crude material that was purified by chromatography on silica gel using ethyl acetate/hexanes to give **54** (0.95 g).

5

Synthesis of 55: 0.6 g of **54** was dissolved in 20 mL dry methylene chloride. The reaction mixture was purged with nitrogen and placed in an ice bath. To this was added, 2 mL of a 2M oxalyl chloride in methylene chloride. The reaction mixture was stirred at room temperature for 1 h and then at room temperature for 2 h. Reaction mixture was concentrated on a rotary and after drying on a vacuum pump for 15 min. the residue (a yellow solid) was dissolved in 15 mL dry methylene chloride, to which was added 2 mL of a 2M solution of dimethylamine in tetrahydrofuran. 30 minutes later the reaction mixture was concentrated and the residue was taken up in ethyl acetate and washed with 10% aq. sodium carbonate, saturated sodium chloride, dried over anhydrous sodium sulfate and filtered. Concentration gives the crude product that was chromatographed on silica gel using ethyl acetate/hexanes to give **55** (0.78 g).

10
15

Synthesis of 57: 7.65g (19.97 mmol) of the indole **56** was dissolved in 80 mL anhydrous dimethylformamide and was cooled in an ice-bath. Under an inert atmosphere, 40 mL (2M solution in dichloromethane) of oxalylchloride was added dropwise over 20 min. After stirring for 15 min. at 0 °C, the reaction was allowed to warm up to ambient temperature and stirring continued for 1h. The reaction mixture was poured in to water and extracted with ethylacetate (3X 100 mL). The combined extract was washed with water and dried

20






over sodium sulfate and evaporated. The product was further purified by column chromatography with ethylacetate:hexane (30:70) to yield 8g (97.5%) of the aldehyde **57**.












Synthesis of 58: To a solution of 1-methylimidazole (149 mg) in tetrahydrofuran (10 mL) was added n-butyl lithium (1.14 mL, 1.82 mmol, 1.6 M in hexane) dropwise at -78 °C. The reaction mixture was stirred at -40 °C for 30 min. **57** (500 mg) in tetrahydrofuran (10 mL) was added. The reaction mixture was warmed to ambient temperature and stirred overnight. The reaction mixture was quenched with ammonium chloride (saturated), extracted with ethyl acetate. The combined organic layer was washed with brine, dried and concentrated. The crude mixture was reflux with phenyl hydrazine (0.6 mL) in ethanol (10 mL) to remove **57**, which was not consumed in the reaction. Removal of the solvent, the residue was purified by chromatography on silica gel eluting with ethyl acetate:hexane (4:1) to give 120 mg of **58**












Using the foregoing procedures, the compounds of Tables 2 and 3 were prepared and many tested for their ability to inhibit p38- α kinase. It was found that the compounds in Tables 2 and 3 provide IC₅₀ values for inhibition of p38- α in the range of 0.1-1.5 μ Mol.











56

Table 2

Compd. #	STRUCTURE	MW (Calcd.)	MW (Obsd.)
1		466	466
2		452	453
3		535	534
4		573	573
5		480	480












6		418	418
7		551	551
8		524	523
9		590	590
10		521	520
11		620	620
12		592	592
13		579	580
14		523	522
15		509	509
16		484	484












17		567	567
18		593	592
19		537	537
20		526	525
21		678	678
22		579	578
23		522	522
24		650	650
25		480	480
26		648	648
27		549	548

28		620	620
29		597	596
30		539	538
31		519	519
32		553	553
33		513	513
34		609	609
35		592	591
36		596	595
37		542	541












38		571	571
39		541	541
40		494	494
41		548	548
42		570	570
43		514	513
44		490	490
45		595	595
46		566	566
47		537	537
48		573	573











49		536	536
50		543	543
51		509	509
52		507	507
53		572	572
54		565	565
55		599	599
56		537	537
57		513	513
58		456	456
59		485	485

60		551	551
61		511	511
62		499	500
63		543	543
64		584	584
65		493	493
66		494	494
67		477	477
68		542	542
69		584	584
70		530	529

71		512	511
72		523	522
73		539	539
74		495	495
75		512	511
76		528	528
77		499	499
78		552	551
79		512	511
80		498	497
81		496	495

82		525	525
83		405	405
84		510	509
85		540	539
86		485	486
87		495	495
88		552	551
89		508	508
90		562	562
91		558	558
92		539	539

93		542	542
94		590	590
95		528	528
96		555	555
97		510	509
98		497	497
99		527	527
100		550	550
101		569	569
102		527	527
103		526	525

104		528	528
105		526	525
106		540	539
107		538	537
108		498	498
109		524	523
110		542	541
111		530	529
112		499	500
113		508	508





















114		542	541
115		504	504
116		492	504













Table 3

Compd. #	MOLSTRUCTURE	MW (Calcd.)	MW (Obs.)
117		472.5858	472.5858
118		404.4636	404.4636
119		390.4368	390.4368
120		502.6116	502.6116
121		558.6752	558.6752

122		458.559	458.559
123		389.4527	389.4527
124		420.4626	420.4626
125		516.6384	516.6384
126		504.6027	504.6027
127		422.4537	422.4537
128		525.021	525.021
129		434.4894	434.4894
130		422.4537	422.4537
131		438.4527	438.4527
132		452.4795	452.4795
133		408.4269	408.4269


134		420.4626	420.4626
135		391.4249	391.4249
136		528.5582	528.5582
137		435.4775	435.4775
138		419.4785	419.4785
139		486.6126	486.6126
140		511.547	511.547
141		507.559	507.559
142		505.5868	505.5868
143		574.6931	574.6931
144		465.5222	465.5222
145		437.4686	437.4686

146		480.9931	480.9931
147		518.6106	518.6106
148		535.0845	535.0845
149		460.5748	460.5748
150		548.6553	548.6553
151		520.6017	520.6017
152		446.548	446.548
153		450.4677	450.4677
154		494.5639	494.5639
155		511.0189	511.0189
156		606.6911	606.6911
157		521.5858	521.5858

158		490.6006	490.6006
159		506.5749	506.5749
160		490.6006	490.6006
161		536.6007	536.6007
162		498.9832	498.9832
163		469.9415	469.9415
164		541.02	541.02
165		511.9783	511.9783
166		497.9951	497.9951
167		497.9951	497.9951
168		483.9683	483.9683
169		539.0478	539.0478


170		549.6434	549.6434
171		476.5738	476.5738
172		476.5738	476.5738
173		476.5738	476.5738
174		469.9415	469.9415
175		479.549	479.549
176		513.01	513.01
177		494.5639	494.5639
178		534.6285	534.6285
179		508.5907	508.5907
180		522.6175	522.6175
181		483.5123	483.5123

73




Claims

1. A compound of the formula:



and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein

 represents a single or double bond;

one Z^2 is CA or CR^8A and the other is CR^1 , CR^1_2 , NR^6 or N wherein each R^1 , R^6 and R^8 is independently hydrogen or noninterfering substituent;

A is $-W_i-COXY_j$ wherein Y is COR^2 or an isostere thereof and R^2 is hydrogen or a noninterfering substituent, each of W and X is a spacer of 2-6Å, and each of i and j is independently 0 or 1;

Z^3 is NR^7 or O;

each R^3 is independently a noninterfering substituent;

n is 0-3;

each of L^1 and L^2 is a linker;

each R^4 is independently a noninterfering substituent;

m is 0-4;

Z^1 is CR^5 or N wherein R^5 is hydrogen or a noninterfering substituent;

each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3;

Ar is an aryl group substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; and

the distance between the atom of Ar linked to L^2 and the center of the α ring is 4.5-24Å.

2. The compound of claim 1 wherein A is $COXjCOR^2$, and

wherein R^2 is H, or is straight or branched chain alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroalkyl, heteroaryl, or heteroarylalkyl, each optionally substituted with halo, alkyl, heteroalkyl, SR, OR, NR_2 , OCOR, NRCOR, $NRCONR_2$, $NRSO_2R$, $NRSO_2NR_2$,

OCOR₂, CN, COOR, CONR₂, COR, or R₃Si wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof, or

wherein R² is OR, NR₂, SR, NRCONR₂, OCONR₂, or NRSO₂NR₂, wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof, and

5 wherein two R attached to the same atom may form a 3-8 member ring and wherein said ring may further be substituted by alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroalkyl, heteroaryl, heteroarylalkyl, each optionally substituted with halo, SR, OR, NR₂, OCOR, NRCOR, NRCONR₂, NRSO₂R, NRSO₂NR₂, OCONR₂, or R₃Si wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof wherein
10 two R attached to the same atom may form a 3-8 member ring, optionally substituted as above defined; and

X, if present, is alkylene.

3. The compound of claim 1 wherein Y is an isostere of COR².

4. The compound of claim 3 wherein Y is tetrazole; 1,2,3-triazole;
15 1,2,4-triazole; or imidazole.

5. The compound of claim 1 wherein each of i and j is 0.

6. The compound of claim 2 wherein j is 0.

7. The compound of claim 1 wherein Z³ is NR⁷.

8. The compound of claim 7 wherein R⁷ is H or is optionally substituted alkyl,
20 alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, or is SOR, SO₂R, RCO, COOR, alkyl-COR, SO₃R, CONR₂, SO₂NR₂, CN, CF₃, NR₂, OR, alkyl-SR, alkyl-SOR, alkyl-SO₂R, alkyl-OCOR, alkyl-COOR, alkyl-CN, alkyl-CONR₂, or R₃Si, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof.

9. The compound of claim 8 wherein R⁷ is H, or is optionally substituted
25 alkyl, or acyl.

10. The compound of claim 1 wherein both k and l are 1.
11. The compound of claim 1 wherein L^1 is CO, CHOH or CH_2 .
12. The compound of claim 11 wherein L^1 is CO.
13. The compound of claim 1 wherein Z^1 is N.
- 5 14. The compound of claim 1 wherein Z^1 is CR^5 wherein R^5 is H, OR, NR_2 , SR or halo, wherein each R is independently H, alkyl, alkenyl or aryl or the heteroatom-containing forms thereof,
- 10 15. The compound of claim 1 wherein L^2 is alkylene (1-4C) or alkenylene (1-4C) optionally substituted with a moiety selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR_2 , SR, SOR, SO_2R , OCOR, NRCOR, $NRCONR_2$, $NRCOOR$, $OCONR_2$, RCO, COOR, alkyl-OOR, SO_3R , $CONR_2$, SO_2NR_2 , $NRSO_2NR_2$, CN, CF_3 , R_3Si , and NO_2 , wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof, and wherein two substituents on L^2 can be joined to
15 form a non-aromatic saturated or unsaturated ring that includes 0-3 heteroatoms which are O, S and/or N and which contains 3 to 8 members or said two substituents can be joined to form a carbonyl moiety or an oxime, oximeether, oximeester or ketal of said carbonyl moiety.
16. The compound of claim 15 wherein L^2 is unsubstituted alkylene.
- 20 17. The compound of claim 15 wherein L^2 is unsubstituted methylene, methylene substituted with alkyl, or $-CH=$.
- 25 18. The compound of claim 1 wherein Ar is optionally substituted with 0-5 substituents selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR_2 , SR, SOR, SO_2R , OCOR, NRCOR, $NRCONR_2$, $NRCOOR$,

5 OCONR₂, RCO, COOR, alkyl-OOR, SO₃R, CONR₂, SO₂NR₂, NRSO₂NR₂, CN, CF₃, R₃Si, and NO₂, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof, and wherein two of said optional substituents on adjacent positions can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members.

19. The compound of claim 18 wherein Ar is optionally substituted phenyl.

20. The compound of claim 19 wherein said optional substitution is by halo, OR, or alkyl.

10 21. The compound of claim 20 wherein said phenyl is unsubstituted or has a single substituent.


22. The compound of claim 1 wherein R⁴ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR₂, SR, SOR, SO₂R, OCOR, NRCOR, NRCONR₂, NRCOOR, OCONR₂, RCO, COOR, alkyl-OOR, SO₃R, 15 CONR₂, SO₂NR₂, NRSO₂NR₂, CN, CF₃, R₃Si, and NO₂, wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof and two of R⁴ on adjacent positions can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members, or R⁴ is =O or an oxime, oximeether, oximeester or ketal thereof.

20 23. The compound of claim 22 wherein each R⁴ is halo, OR, or alkyl.

24. The compound of claim 23 wherein m is 0, 1, or 2.

25. The compound of claim 24 wherein m is 2 and both R⁴ are alkyl.


26. The compound of claim 1 wherein each R³ is halo, alkyl, heteroalkyl, OCOR, OR, NRCOR, SR, or NR₂, wherein R is H, alkyl, aryl, or heteroforms thereof.

27. The compound of claim 26 wherein R^3 is halo or alkoxy.
28. The compound of claim 27 wherein n is 0, 1 or 2.
29. The compound of claim 1 wherein L^1 is coupled to the α ring at the 4-, 5- or 6-position.
- 5 30. The compound of claim 1 wherein Z^2 at position 3 is CA or CH^1A .
31. The compound of claim 30 wherein the Z^2 at position 2 is CR^1 or CR^1_2 .
32. The compound of claim 31 wherein R^1 is hydrogen, or is alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, NH-aroyl, halo, OR, NR_2 , SR, SOR, SO_2R , OCOR, NRCOR, $NRCONR_2$,
10 $NRCOOR$, $OCONR_2$, RCO, COOR, alkyl-OOR, SO_3R , $CONR_2$, SO_2NR_2 , $NRSO_2NR_2$, CN, CF_3 , R_3Si , and NO_2 , wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof and two of R^1 can be joined to form a fused, optionally substituted aromatic or nonaromatic, saturated or unsaturated ring which contains 3-8 members.
33. The compound of claim 32 wherein each R^1 is selected from the group
15 consisting of H, alkyl, acyl, aryl, arylalkyl, heteroalkyl, heteroaryl, halo, OR, NR_2 , SR, NRCOR, alkyl-OOR, RCO, COOR, and CN, wherein each R is independently H, alkyl, or aryl or heteroforms thereof.
34. The compound of claim 30 wherein Z^2 at position 2 is N or NR^6 .
35. The compound of claim 34 wherein R^6 is H, or alkyl, alkenyl, alkynyl, aryl, arylalkyl, acyl, aroyl, heteroaryl, heteroalkyl, heteroalkenyl, heteroalkynyl, heteroalkylaryl, or is SOR, SO_2R , RCO, COOR, alkyl-COR, SO_3R , $CONR_2$, SO_2NR_2 , CN, CF_3 , or R_3Si wherein each R is independently H, alkyl, alkenyl or aryl or heteroforms thereof.
- 20 36. The compound of claim 1 wherein  represents a double bond.

37. The compound of claim 1 wherein the distance between the atom on Ar linked to L² and the center of the α ring is 7.5-11 Å.

38. The compound of claim 1 wherein the compound of formula (1) is selected from the group consisting of compounds shown in Tables 2 and 3 herein.

5 39. A pharmaceutical composition for treating conditions characterized by enhanced p38- α activity which composition comprises a therapeutically effective amount of a compound of the formula



10 and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein

represents a single or double bond;

one Z² is CA or CR⁸A and the other is CR¹, CR₂¹, NR⁶ or N wherein each R¹, R⁶ and R⁸ is independently hydrogen or noninterfering substituent;

15 A is -W_i-COX_jY wherein Y is COR² or an isostere thereof and R² is hydrogen or a noninterfering substituent, each of W and X is a spacer of 2-6 Å, and each of i and j is independently 0 or 1;

Z³ is NR⁷ or O;

each R³ is independently a noninterfering substituent;

n is 0-3;

20 each of L¹ and L² is a linker;

each R⁴ is independently a noninterfering substituent;

m is 0-4;

Z¹ is CR⁵ or N wherein R⁵ is hydrogen or a noninterfering substituent;

each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3;


Ar is an aryl group substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; and

the distance between the atom of Ar linked to L^2 and the center of the α ring is 4.5-24Å.

40. The composition of claim 39 which further contains an additional therapeutic agent.

41. The composition of claim 40 wherein said additional therapeutic agent is a corticosteroid, a monoclonal antibody, or an inhibitor of cell division.

42. A method to treat a condition mediated by p38- α kinase comprising administering to a subject in need of such treatment a compound of the formula:



and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein

represents a single or double bond;

one Z^2 is CA or CR^8A and the other is CR^1 , CR^1_2 , NR^6 or N wherein each R^1 , R^6 and R^8 is independently hydrogen or noninterfering substituent;

A is $-W_i-CO-X_jY$ wherein Y is COR^2 or an isostere thereof and R^2 is hydrogen or a noninterfering substituent, each of W and X is a spacer of 2-6Å, and each of i and j is independently 0 or 1;

Z^3 is NR^7 or O;

each R^3 is independently a noninterfering substituent;

n is 0-3;

each of L^1 and L^2 is a linker;

each R^4 is independently a noninterfering substituent;

m is 0-4;

Z^1 is CR^5 or N wherein R^5 is hydrogen or a noninterfering substituent;

each of l and k is an integer from 0-2 wherein the sum of l and k is 0-3;

Ar is an aryl group substituted with 0-5 noninterfering substituents, wherein two noninterfering substituents can form a fused ring; and

5 the distance between the atom of Ar linked to L² and the center of the α ring is 4.5-24Å.

43. The method of claim 42 wherein said condition is a proinflammation response.

44. The method of claim 43 wherein said proinflammation response is multiple sclerosis, IBD, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis,
10 other arthritic conditions, sepsis, septic shock, endotoxic shock, Gram-negative sepsis, toxic shock syndrome, asthma, adult respiratory distress syndrome, stroke, reperfusion injury, CNS injury, psoriasis, restenosis, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, a bone resorption disease, graft-versus-host reaction, Crohn's Disease, ulcerative colitis, Alzheimer's or pyresis.

INTERNATIONAL SEARCH REPORT

Int lonal Application No

PCT/US 00/14003

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/06 A61K31/404 A61P43/00 C07D209/24 C07D401/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 06715 A (SMITHKLINE BEECHAM CORPORATION) 19 February 1998 (1998-02-19) cited in the application claims -----	1, 39
A	WO 98 28292 A (SMITHKLINE BEECHAM CORPORATION) 2 July 1998 (1998-07-02) claims -----	1, 39
A	WO 99 00357 A (VERTEX PHARMACEUTICALS INCORPORATED) 7 January 1999 (1999-01-07) page 3 -page 4 -----	1, 39



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 October 2000

Date of mailing of the international search report

25/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/14003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9806715 A	19-02-1998	EP 0922042 A	16-06-1999
W0 9828292 A	02-07-1998	NONE	
W0 9900357 A	07-01-1999	US 6093742 A	25-07-2000
		AU 8377698 A	19-01-1999
		EP 0993441 A	19-04-2000